

PROCEEDINGS OF THE EIGHTH MEETING OF

THE COMMITTEE ON FOREST TREE BREEDING

IN CANADA

Held at the University of New Brunswick, Fredericton, N.B., August 16-18, 1962.

PART II

MEMBERS ' PROGRESS REPORTS

Assembled and Distributed by the Forest Research Branch, Department of Forestry, Ottawa.

Part I, Minutes and Discussions, received restricted distribution to Committee members only. Part II received wider distribution to persons and organizations actively engaged or interested in forest tree breeding and improvement.

CONTENTS

1	LIST OF ACTIVE MEMBERS OF THE COMMITTEE	<u>Pa</u>	<u>ge No.</u>
⊥•	ON FOREST TREE BREEDING IN CANADA, SEPTEMBER 1962		2
2.	MEMBERS ' PROGRESS REPORTS		-
	M. G. Boyer	A	1-3
	A. J. Carmichael	В	1-5
	L. P. Chiasson	С	1-2
	W. H. Cram	D	1-11
	B. W. Dance	Е	1-3
	L. F. Ebell	F	1-2
	J. L. Farrar	G	1
	D. P. Fowler	H	1-4
	D. A. Fraser	I	1-2
	W. V. Hancock	K	1-3
	C. Heimburger	L	1-6
	M. J. Holst	М	1-25
	H. G. MacGillivray	0	1-14
	R. J. Moore	P	1-2
	A. L. Orr-Ewing	Q	1-3
	L. Parrot	R	1-2
	C. R. Sullivan	U	1-3
	0. Sziklai	V	1-7

1.	ΓI	ST OF	F ACTIVE MEMBI BREEDING IN CA	ERS OF THE COMMITTEE ON FOREST TREE ANADA, SEPTEMBER 1962
Dr.	М.	G. I	Boyer	Canada Dept. of Forestry, Forest Entomology and Pathology Branch, Sillery, Que.
Dr.	F.	C. H	Bent	Acadia University, Wolfville, N.S.
Mr.	A.	J. (Carmichael	Ontario Dept. of Lands and Forests, Research Branch, Maple, Ont.
Dr.	L.	P. (Chiasson	St. Francis Xavier University, Antigonish, N.S.
Dr.	W.	н. с	ram	Canada Dept. of Agriculture, Research Branch, Indian Head, Sask.
Mr.	в.	W. [ance	Canada Dept. of Forestry, Forest Entomology and Pathology Branch, Maple, Ont.
Mr.	J.	T. D	Oorland	Kimberly-Clark Pulp and Paper Co. Ltd., Longlac, Ont.
Mr.	W.	G. D	yer	Ontario Dept. of Lands and Forests, Timber Branch, Toronto, Ont.
Dr.	L.	F.E	bell	Canada Dept. of Forestry, Forest Research Branch, Victoria, B.C.
Dr.	J.	L.F	arrar	University of Toronto, Faculty of Forestry, Toronto, Ont.
Mr.	D.	P.F	owler	Ontario Dept. of Lands and Forests, Research Branch, Maple, Ont.
Dr.	D.	A.F	raser	Canada Dept. of Forestry, Forest Research Branch, Chalk River, Ont.
Mr.	с.	Gagn	on	Canada Dept. of Forestry, Forest Entomology and Pathology Branch, Sillery, Que.
Mr.	W.	₩. Н	ancock	Canada Dept. of Forestry, Forest Products Research Branch, Vancouver, B.C.
Mr.	J.	С. Н	eaman	British Columbia Forest Service, Research Division, Victoria, B.C.
Dr.	c.	С. Н	eimburger	Ontario Dept. of Lands and Forests, Research Branch, Maple, Ont.

1

- 2 -

	Mr.	M. J. Holst (Chairman)	Canada Dept. of Forestry, Forest Research Branch, Chalk River, Ont.
	Dr.	A. H. Hutchinson	University of British Columbia, Dept. of Biology and Botany, Vancouver, B.C.
	Mr.	H. G. MacGillivray	Canada Dept. of Forestry, Forest Research Branch, Fredericton, N.B.
	Dr.	R. J. Moore	Canada Dept. of Agriculture, Plant Research Institute, Ottawa, Ont.
	Mr.	E. K. Morgenstern (Secretary)	Canada Dept. of Forestry, Forest Research Branch, Chalk River, Ont.
	Dr.	A. L. Orr-Ewing	British Columbia Forest Service, Research Division, Victoria, B.C.
	Dr.	L. Parrot	Université Laval, Faculté d'Arpentage et de Génie Forestier, Québec, Qué.
	Dr.	P. J. Pointing	Canada Dept. of Forestry, Forest Entomology and Pathology Branch, Sault Ste. Marie, Ont.
c	Mr.	0. Sziklai	University of British Columbia, Faculty of Forestry, Vancouver, B.C.
,	Mr.	C. W. Yeatman	Canada Dept. of Forestry, Forest Research Branch, Chalk River, Ont.

- 3 -

Report to Committee on Forest Tree Breeding

1962

Michael G. Boyer

Canada Department of Forestry Maple, Ontario.

One request by tree breeders with respect to white pine blister rust work has been for methods of selecting seedlings and grafted material which would establish with certainty the resistance or susceptibility of the host. While probably the most effective method would involve the characterization of the rust resistant factors, the aspect we are at present investigating, techniques of inoculation could perhaps be improved. During initial studies on the physiology of the disease some techniques of inoculation were investigated to provide information on specific aspects of the disease. Four were attempted on white pine seedlings in the laboratory; a) pre-germinated teliospores prepared by removing telia from <u>Ribes</u> leaves incubating on 2 per cent water agar at 16° C for 12 hours and applying them to parts of the seedling, b) acciospore inoculation of leaves and stems with and without wounding and with or without <u>Ribes</u> leaf extract, c) implantation of diseased leaf tissue in the hypocotyl region of the seedling and, d) inoculation with excised <u>Ribes</u> leaves.

Infection was attempted by incubating seedlings at 15°C in the dark for five days in closed containers. Number of inoculations and successful transmission are indicated in Table I. A brief discussion of each method follows.

Table I

Number of Seedlings	Method	Place of Inoculation	Number	Number
50	aeciospores*	hypocotyl	50	0
6	1¥ × ×	cotyledons	20	0
10	11	primary leaves	23	0
}	excised telia	cotyledons	122	2
66)	11	primary leaves	120	5
~~}	99 · ·	true leaves	100	4
)	f7	hypocotyl	21	0
34	infected white pine leaves	hypocotyl	34	3.
24	Ribes leaves			24

Effect of method of inoculation on infection of white pine seedlings

Methods of acciospore inoculation are not listed completely because none was successful.

Inoculation with acciospores

Scholz's report^{*} that aeciospores caused canker-like infections when wound inoculated on white pine stems suggested a possible simple method of infecting seedlings. As a consequence seedlings were inoculated at various locations (Table I) either directly without wounding or by brushing viable aeciospores into a small longitudinal wound made with a sharp scalpel. An extract of <u>Ribes</u> leaves was included in some inoculations. To complement this study field inoculations were carried out on 50 first, second and third year shoots of approximately 15-year-old white pine. Under these circumstances no infection developed after five months although germination of aeciospores in the wounds were observed in all embedded material. Since in the case of seedlings infection could be induced on all parts by other methods it would appear that infection may be attained only under more restricted circumstances,

Inoculation with excised telia

The purpose of using excised telia was two-fold. By selections of vigorous productive telial columns it was hoped to be able to assure infection at any point and also to study the effect of distance of inoculum from the stem. Since viable telial columns were selected and transfer after 12 hours to 2 per cent water agar resulted in the active production of sporidia the failure to obtain a higher incidence of infection is puzzling. Penetration of sporidia on hypocotyls was observed frequently. Further work has been limited by space, and modification of the method has not been attempted as yet. However, it has been possible to show that all of the leaves are susceptible and while no infections on the hypocotyl were observed the fact that sporidial penetration was evident suggested that some resistance to sporidial infection is present.

Inoculation with the diseased pine leaf sections

Attempts to inoculate seedlings by transfer of thin sections of diseased leaf tissue into the pith region at the base of the cotyledonary node was prompted by the observation that the transfusion tissue of first year leaves promptly forms callus when wounded and placed in a high humidity. The possibility of grafting infected tissue to seedlings was considered. The technique involved the transfer of approximately one mm. leaf sections in which the fungus was located in the transfusions tissue into the base of the node by making a longitudinal incision and inserting the leaf section in such a way as to obtain contact between the pith cells of the hypocotyl and the transfusion tissue of the leaf. In two of the three successful transfers examined in detail the fungus had apparently moved directly into the proliferating cells of the pith and from thence into the ray cells of the xylem. Of the 34 inoculations (Table I)

الم الم الم الم

and an end of the state of a state of a state of the stat

¥

See Scholz, E. 1960. Zuchter 30(2), 61-72.

made at about 2 weeks after germination (the cotyledonary stage) none died and further growth impediment was slight or not observable. Failure to obtain a higher incidence of infection was investigated with free-hand material. In most cases the inability to obtain contact between the tissues either because of improper placement of leaf sections or because the leaf sections were not cut at right angles appeared to be the primary cause of failure.

Inoculation with excised leaves of Ribes

This method which has been used frequently in the past was employed to determine if the conditions used for inoculation, i.e., 5 days at 15° C in the dark, were satisfactory. Of the 24 seedlings inoculated 100 per cent exhibited dense multiple infections within one month.

WHEN REPLYING KINDLY QUOTE THIS FILE NUMBER

Reforestation



DEPARTMENT OF LANDS AND FORESTS

Toronto, Ontario

REPORT TO COMMITTEE

ON

FOREST TREE BREEDING

August, 1962

Hon. J. W. Spooner Minister F. A. MacDougall Deputy Minister

J. A. Brodie, Chief Timber Branch REPORT TO COMMITTEE ON FOREST TREE BREEDING

August 1962

A. J. Carmichael

Ontario Department of Lands and Forests

1 **PROVENANCE TESTS**

A. Red Pine

A red pine provenance test composed of 12 Ontario seed sources was outplanted in 1956. Measurements of total height, after 10 growing seasons from seed, were made in fall 1961 at Cochrane, Sault Ste. Marie and Lindsay Districts in northern, central and southern Ontario.

Mean Height in Feet for Red Pine Sources

Provena	ince	Planting	Location -	Site Region
Site Region	Locality	3E	5E	6E
3W 4S 4E 5E	Port Arthur Eagle River Timagami Barry's Bay Douglas Golden Lake Mattawa	1.6 1.5 1.7 1.5 1.7 1.7	3.1 2.9 3.2 3.2 3.3 3.2 3.3	3.9 3.8 3.9 4.0 4.1 3.7 2.7
6E	Pembroke Thessalon Average Angus Bruce Kemptville Average	1.7 1.6 1.61 1.6 1.7 1.7 1.67	3.3 3.20 3.1 3.2 3.4 3.25	4.0 3.6 3.86 3.9 3.8 4.2 3.97
General Av	erage	1.62	3.18	3.90

An analysis of variance was completed following the design prepared by Dr. D. B. DeLury. This indicated, first, that the experiment had failed to show any significant difference between the heights of different provenances at each of the planting sites; and second, that there was a highly significant difference between the heights of all provenances when grown in different Site Regions. The height of all provenances was greater in the southern planting.

B-1

Source	<u>df</u>	<u>55</u>	ms	F	F.05	
Bet. Climatic Regions Reps. (Within Regions)	2 9	129.5 3.3	64.7 0.37	176.4	4.3 **	
Bet. Provenances Regions x Provenances Reps. (Within Regions)x	11 22	1.6 0.93	0.14 0.04	41 41	2.45 1.84	
Provenances	<u>99</u> 143	17.1	0.17			

SULTIARY OF ANALYSIS

B. Jack Pine

A jack pine provenance test composed of 12 Ontario seed sources was outplanted in 1955. Measurements of total height after 10 growing seasons from seed, were made in fall 1961, at Swastika and Pembroke Districts in northern and central Ontario.

Mean Height in Feet for Jack Pine Sources

	Provena	ince	Planting	Location .	- Site Region
Site	Region	Locality	4E	5E	Average
	3W	Geraldton Kab Lake Pickle Lake	5.8 7.0	5.5 6.2 5.8	5.65 6.60 6.00
3E Co T: F:		Connaught Timmins Franz	6.6 7.1 6.1 5.6	6.2 7.1 6.0	6.40 7.10 6.05 5.95
	4S	Sioux Lookout	7.3	6.6	6.95
	4W	Goldie Hardwick	6.ඊ 6.9	7•2 6•⊃	7.00 6.85
	5E	Markstay Algoma	7.4 6.4	7.2 6.4	7.30 6.40
	General A	verage	6.40	6.60	· · · · ·

An analysis of variance was completed following the design prepared by Dr. D. B. DeLury. This indicated, first, that the experiment had failed to show any significant difference between the height of jack pine grown in different site regions; and second, that there was a highly significant difference in the height growth of different provenances, when the region in which they were grown was not considered.

Source	$\underline{\mathtt{Df}}$	SS	ms	F	F.05
Bet. Climatic Regions Reps. (Within Regions)	1 6	1.16 2.15	1.16 0.35	3.24	5.99
Bet. Provenances Regions x Provenances Reps (Within Regions)r	11 11	27.84 4.60	2.53 0.42	4.71 ∢1	1.94** 1.94
Provenances	<u>66</u> 95	33.80	0.51		

Summary of Analysis

11 SEED ORCHARDS

The emphasis on seed orchard establishment has been reduced currently in favour of the development of seed production areas. The present target is for 10,000 grafts to be carried out annually at five nursery locations (Sb 3000, Sw 3000, Pr 25000, Pw 1500).

Some changes have been made in grafting procedure to improve scion development. Grafts made at an earlier stage of stock plant development, seem to be beneficial for both spruce and red pine. Spruce are grafted when stock plant root activity is evident and before bud swelling occurs. Red pine are grafted when bud swelling is evident but beforesshoot elongation has progressed.

The release of developing spruce shoots from constriction within bud scales has been necessary and has resulted in greater survival with better shoot development.

Various methods for fall and early winter storage of understock have been employed to simplify maintenance procedures. Storage of stock growing in 5 inch clay pots has been carried out adequately in unheated sheds with limited natural light. Red pine and white pine have been carried over in excellent condition when no mulch was used on the pots. Some difficulties were experienced when pots and trees were covered with a conifer brush mulch. Further testing is indicated to obtain uniform results and these are being developed at each nursery.

The selection of orchard planting sites for spruce and pine has led to the examination of hardwood areas for the exclusion of foreign pollen. A few sites are available, however the difficulty in locating accessible areas that can be cleared at a reasonable cost, has indicated the possible value of using locations on large burned areas and controlling future planting around the orchard site.

111 SEED PRODUCTION AREAS

A. <u>General</u>

The selection of stands suitable for seed production area development has been hampered by the apparent lack of younger age classes. Therefore, older stands are being selected in which there are from 50 to 100 seed producing trees per acre at heights of from 40 feet to 80 feet. Cone collection from such trees will require the use of tree-climbing bicycles and thus increase seed collection costs. The use of truck or tractor-mounted ladders may be possible in some locations, however, stands of better quality are not always readily accessible and will require less cumbersome climbing equipment.

When tree-climbing bicycles are used extensively, it will be possible to train staff and develop collecting techniques that should enable more extensive collections from the better stands.

The development of plantations from selected seed sources offers one solution to the problem. Some white and black spruce sources are being grown for this purpose. A more extensive local development of select plantations, using seed from the best stands in the district, would provide ultimately for a cone collection source to supplement that from seed orchards and seed production areas in thinned natural stands.

B. <u>White Spruce - Site Region 3W</u>

A location at Pagwa in Geraldton District was scarified to induce seedlings beneath an excellent white spruce stand. Regeneration on the 20 acre block resulted in 20,000 seedlings per acre at the end of the first year. Subsequent counts have not been made, but there are still an adequate number of 2 year old plants.

C. <u>Red Pine, Lynn Tract, Simcoe County</u>

Trees of 12 ft. to 20 ft. in height were thinned to a 12 ft. x 12 ft. and an 18 ft. x 18 ft. spacing in 1957. Five years later the crowns of trees at the 12 ft. spacing again required release, whereas the 18 ft. spacing will be adequate for at least another 5 years and possibly 10 years. The 18 ft. spacing has been accepted generally for thinning in stands up to a 40 ft. height.

The seasonal changes in foliar nutrient levels are being examined on this area by Professor K. Armson. His work will determine the best sampling procedure to assess the nutrient capital of young red pine. When coupled with a measure of the tree's photosynthetic potential it should be possible to differentiate those trees that produce cone crops above their normal level and thus evaluate the true effects of fertilizer treatments. The physiological factors involved in flower induction and the effects of fertilizer applications on these still need to be examined.

D. <u>Wood Quality</u>

The development of a black spruce seed production area from seed obtained at Parnell Township in Kapuskasing District, necessitated the felling of 32 trees for cone collection. Each of the felled trees was sectioned and blocks 1 foot in length were taken to represent each 100 inch level above the ground. Blocks were measured to develop stem analyses for each tree and then shipped to the Forest Products Laboratory in Ottawa for cellular examination.

REPORT ON TREE BREEDING - 1962

L. P. Chiasson St. Francis Xavier University Antigonish, N. S.

After the loss of promising hybrid material in 1960, efforts were continued to produce interspecific Abies hybrids.

In the spring of 1960, the following exotic pollens were used on Abies balsamea female cones: A. cephalonica, A. concolor, A. homolepis, A. homolepis var. umbellata, A. koreana, A. lasiocarpa, and A. nordmanniana.

The hybrid seeds were collected in early September, 1960 and stored in a dry room over winter. They were stratified for a month at approximately 5°C, and then sown in specially prepared outdoor beds. These beds were covered with a double layer of cheesecloth, and were sprinkled once a day through the summer months.

Table I shows the results in terms of cones collected for each cross.

	TABLE 1	
Putative Male Parent	No. of cones collected	Seeds germinated-1961
A. cephalonica A. concolor A. homolepis A. homolepis A. homolepis var. umbellata A. koreana A. lasiocarpa A. nordmanniana	10 17 303 84 167 70 68	x x x

The germination was extremely low not only for these potentially hybrid seeds but also for <u>balsamea</u> seeds that had been collected, planted and stored under identical conditions.

In the spring of 1962 the <u>balsamea-lasiocarpa</u> crosses seemed to be the only ones that had overwintered successfully, but more careful examination of the beds is required before a final assessment can be made.

In the spring of 1961, the only exotic pollen available was from A. nordmanniana. A crop of 49 cones presumably containing hybrid seeds were collected, and after stratification, have been planted this spring.

In the spring of 1962, the following exotic pollens have been used on balsamea ovulate cones: alba, cephalonica, cilicica, concolor, ernesti, Fraseri, homolepis, lasiocarpa var. arizonica, nephrolepis, nordmanniana, sachilanensis, sibirica, Veitchii. Only 92 ovulate cones were isolated this spring, as there was a scarcity of these and many of those that were visible were not available for isolation with the equipment at our disposal.

Investigation of the comparative cytology of Abies species is continuing in anticipation of obtaining confirmed interspecific crosses.





h. te s

2 ./

والمتعادين والمراجع . . .

1960-61 Biennial Progress Report

Shelterbelt Tree Breeding

W.H. Cram, Superintendent Forest Nursery Station, Canada Department of Agriculture, Indian Head, Saskatchewan.

GENERAL

Drought conditions, which have prevailed since 1957, were unfavourable to the Nursery and Breeding research programs in 1960 and 1961. 1961 was the driest year on record for this station with an annual precipitation of only 7.5 inches. The frost free period of 1960 was 117 days (May 17 to September 10) and of 1961 was 112 days (May 13 to September 2) and the precipitations in these periods were 4.9" and 1.6" respectively. Heavy losses, due to drought and extreme temperatures, were noted for germination capacity, seedling stand and even mature stands of trees in permanent plantings.

Staff changes included the appointment of Dr. R. Grover in October 1960 as Plant Physiologist, to conduct Herbicidal and seed technology investigations related to tree production, and Mr. P.J. Salisbury, Pathologist in April 1960 to conduct investigations in disease problems related to tree production.

CARAGANA BREEDING

Evaluation of combining ability for self-incompatible selections continued to be the major phase of the Breeding program for <u>Caragana arborescens</u> Lam. Preliminary research led to the selection of vigorous self-incompatible plants from Indian Head material. These selections were cross-incompatible and therefore unsuited for planting in isolated natural-crossing blocks to mass produce intra-specific crosses. This demonstrated the need for new accession material to provide additional self-incompatible plants of genetic diversity and crosscompatible with existing material. Cross-compatibility of nine self-incompatible selections from accession material with pollen of five similar selections from Indian Head material was determined in 1961, with the results listed in Table 1.

Accessi	on	Indian Head Selections (3)						
Selection	Source	Al	V16	V17	B24	₩2		
· (♀)	(country)	(%)	(%)	(%)	(%)	(%)		
D10-13	Sweden	0	0	0	0	0		
D11-8	U,S.A.	0	0	0	0	0		
D14-71	Czech.	0	0	0	0	0		
D16-3	France	0	0	0	0	0		
D16-10	France	0	0	0	0	6		
D14-26	Czech.	0	0	0	0	19		
D14-13	Czech.	0	2 ?	0	0	50		
D14-112	Czech.	0	0	0	17	48		
D14-12	Czech.	0	0	11	15	17		

Table 1. Cross-Compatibility¹ of 9 Self-incompatible Accession Selections with 5 Self-Incompatible Local Selections of Caragana

¹Compatibility as % of cross pollinated flowers setting pods ?Extreme deviate may be contamination due to protruding flowers

The nine selections listed in Table 1 were made in 1959 and 1960, by selfing 140 vigorous plants produced from eight seed accessions. Regretably the 1961 breeding program was seriously disrupted and the results distorted by the prevailing drought and high temperatures. Nevertheless, cross-compatibilities listed in Table 1 suggest four of the accession selections carry the same incompatibility factors as five local selections, and are of the genotype $S_1 S_2$. However, five of the accessions appear to carry one or more different incompatible factors, which permit cross-compatibility with pollen from three local selections. These accessions offer promise as parental material for the establishment of natural-crossing blocks. In the next few years, it will be necessary to repeat and confirm these 1961 compatibility tests, propagate the selections by cuttings and evaluate the vigor potential of hybrid progenies for all possible combinations.

Relative vigor of hybrid caragana progenies, when evaluated in terms of plant height, was found to change until the seedlings reached sexual maturity after 6 years of growth. Height data, as recorded for eleven polycross progenies after 3, 4, 5 and 6 years of growth from seed, have been summarized in Table 2. Relative vigor, as portrayed by the juvenile seedlings after 3 and 4 years of growth, was maintained beyond sexual maturity for only one of the eleven progenies, namely for V-13. On the other hand that of all other progenies changed in the 5th and 6th years of growth. This change was slight or gradual for some progenies but extreme for others. For example the vigor rankings for the polycrosses V-2 and 21-28 changed from fourth and first place to second and third place, respectively, in the 5th and 6th year, whereas that of 21-31 changed from eleventh to first place. As a result ratings for the vigor potential of hybrid caragana progenies should be postponed until the seedlings attain sexual maturity after 6 years' of growth.

Table 2.	Comparative	Mean	Heights	and	Rankings	of	Heights ¹	for	Eleven	Polycross
	Progenies a	s 3,	4, 5, an	d 6	year-old	see	dlings.			

Polycross	Hei	ght in cer	ntimeters		Ran	5		
progeny	3Yr.	4Yr.	5Yr.	6Yr.	3Yr.	41r.	5Yr.	6Yr.
21-31	54	80	145	194	11	11	2	1
V-2	73	109	143	187	1	l	4	2
21-28	72	109	148	184	2	2	i	3
21-16	64	98	140	181	7	8	6	4
V-15	68	97	136	178	6	9	8	5
A-1	69	102	142	177	3	3	5	6
A-3	59	99	134	176	9	7	9	7
21-27	68	100	144	173	5	6	ġ.	8
V - 16	68	101	140	171	4	5	7	9
V - 13	59	92	130	166	10	10	10	10
21-32	64	102	114	156	8	4	11	11

¹Mean height of some 40 seedlings for each year and progeny

Pollen viability and storage investigations were continued for caragana in 1961. One study involved the influence of sucrose content of water-agar culture media on pollen germination. Pollen from two selections of caragana (V16 and V17) was incubated in petri dishes on 2% water-agar media with 1, 5 and 10% sucrose content. Germination was recorded after 5, 20, 52 and 76 hours of incubation with the results given in Table 3. Germination of pollen increased with each increment

Tab.	_e	3.	Pollen	Viability ¹	for	Two	Caragana	Se]	Lections	at	Three	Sucrose	Leve]	Ls
														_

Incubation	_	A-2			V-17			
Period ²	1%	5%	10%	1%	5%	10%	Means	
(Hrs_{\bullet})	(%)	(%)	(%)	(%)	(%)	(%)	· · · · · · · · · · · · · · · · · · ·	
5	45	42	56	32	40	70	47.4	
20	26	40	69	45	56	77	52.1	
52	26	53	63	24	63	88	52.6	
76	43	41	65	33	53	88	53.6	

¹Viability as % germination in one field across petri dish

²Then sample removed and stained with Aceto-carmine for recording germination

in the sucrose content of the agar media for both selections and for all periods of incubation. The data showed that germination after 20 hours of incubation on 2% water-agar substrate with 10% sucrose provides a reliable measure of pollen viability for caragana.

Viability of pollen from bud and bloom stages of caragana after two types of storage at two temperatures was investigated. Buds in full color and freshlyopened flowers were harvested from one selection (V17) on June 1st. These were placed in manila paper and polyethylene bags, then stored at 0° and 40°F. Pollen samples were taken from these buds and open flowers after 7, 13, 28 and 42 days of storage, and the germination recorded after incubation for 24 hours at 70°F on 5% sucrose-agar medium. The following conclusions on viability of caragana pollen were drawn from the results of this study: -

(1) Storage temperature at 40°F was superior to that of 0°F;

(2) Storage in paper bags was better than storage in poly bags;

(3) Pollen viability decreases rapidly from 66% to 8% after 13 days of storage;

(4) Buds in full color if stored in paper bags at 40°F seem to provide the best pollen for extending the period for cross-pollinations in a caragana breeding program.

Four_types_of media were_investigated for the rooting of softwood caragana cuttings in 1960. In all cases the rooting media were placed in wooden flats and cultured on heating cables under intermittent mist. Shoots were collected from one clone at 5 bi-weekly intervals from June 8 to August 3 and made into 4" terminal and basal cuttings. Ten cuttings of each sample of 20 cuttings were dipped in 'Rootone' hormone just prior to planting. Records for rooting were taken 21 days after planting and are summarized in Table 4. Rooting was greatest (38%) in a pure medium of local screened sand, and next highest in Flint-shot Ottawa-silica sand. Peatmoss additives had no beneficial influence, whereas applications of Rootone hormone materially increased rooting.

with (+) and	without	t (-) Hor	mone Treatment	;	_
Rooting	Term	inal	Basa	1	Media
media	+		+	-	Neene
Local screened sand	26	4	72	52	38.5
Ottawa sand Flint shot	2	6	74	52	33.5
Local sand & moss	28	4	44	40	29.0
Ottawa sand & moss	10	16	46	28	24.5
Means	16 -	7	59	43	

Table 4. Rooting¹ for Terminal and Basal Cuttings of Caragana in Four Media

Mean value for 10 cuttings & 5 dates as % of number planted.

SPRUCE BREEDING PROGRAM

Self- and open-compatibility was determined for 31 selections of Colorado spruce (Picea pungens) in 1960. Self-compatibility of these trees ranged from 0 to 46 seeds-per-cone and averaged 12; while open-compatibility ranged from 0 to 118 and averaged 51 seeds-per-cone. This wide range for self-compatibility in the species suggests the potential use of self-incompatible lines for the breeding program.

selections of Colorado spruce, which had differed as to self-compatibility in 1959. Two previous selections, RCl and RC2, which were outstanding for 'blue' needle color, were utilized as pollen parents to test relative cross-compatibilities and subsequent inheritance of needle color. Colorado spruce produced few flowers in 1960 so that it was impossible to conduct all the desired pollinations. The results for 1959 and 1960 appear in Table 5.

	Colorado Spruc	e in 1959 a	nd in 1960)			
	Open-p	ollination	Cross	3- 1960	Self-pollination		
Seedtree	1959	1960	RC-1	RC-2	1959	1960	
	(%)	(%)	(%)	(%)	(%)	(%)	
NC4-5	87.4	50.0	0	0	1.4	0	
NC5-23	65.3	41.5	8.0	14.6	3.5	0	
NC 5-25	128.0	67.0	14.5	34.0	0,6	0	
NC4-51	29.4	15.5	37.0	÷.	1.5	0	
NC5-28	144.8	-	42.7	4.6	32.4	0	
NC4-18	33.3	29.5	0	25.0	0.0	۲ ۸	
NC4-60	61.4	53.8	13.2	15.5	0.0	2.5	
NC5-16	100.6	75.6	10.8	43.8	4.5	6.8	
NC5-58	44.0	28.3	33.8	45.0	1.2		
Weighted N	leans 68.5	51.6-	15.4	26.1		1.5	

Table 5. Open-, Cross-, and Self-compatibility¹ for Nine Selections of

¹Compatibility expressed in terms of the sound seeds-per-cone ²Seed set reduced by losses due to pre-harvest shedding of seed *No male flowers in 1960, - - no data available.

Seed set was generally less in 1960 than in 1959. For example the average set from open-pollination in 1960 was 45 seeds-per-cone, only 66% of that for the same trees in 1959. Nevertheless, the efficiency of the techniques for artificial cross-pollinations approached 40% of that for natural open-pollinations, despite possible cross-incompatibility interactions. Differential meed sets for pollen of RC1 and RC2 with the seedtrees NC5-28 and NC4-18 suggest that degrees of crosscompatibilities exist within the species. Failure of both pollens to effect seedset with the seedtree NC4-5 indicates the possibility that cross-incompatibilities exist within the species. However, inconsistency of the seedsets following selfpollination of six trees in 1959 and 1960 demonstrates a need for further research to ensure comparable results from year to year.

<u>Self-compatibility</u> was determined for 41 selections of Colorado spruce in 1961 and in addition self- and cross-compatibility were checked for 10 selections showing low self-compatibility in 1960. Regretably, drought, heat and seed insect problems seriously disrupted the 1961 spruce breeding program and generally reduced seed set. These results suggested precautions must be taken in the future to reduce insects and temperatures within isolation bags.

Survival, height and needle coloration data were recorded in 1961 for 13-year-old plants in open-pollination progenies of 24 Colorado spruce selections. Similar data were recorded for these plants in 1953, and height data were compiled in 1956. The results have been listed in Table 6. An average survival of 91% demonstrates the ability of Colorado spruce, which had become adequately established under favorable conditions from 1954 to 1956, to withstand subsequent adverse conditions while in the juvenile stage. Blue coloration of the needles of 13-year-old plants bore no relationship to that of the same plants when 5 years old. All plants of three progenies exhibited some degree of blue needle color, and only three progenies had high average color ratings exceeding the mean. Height of plants after 13 years of growth was closely related to that after 8 years of growth as indicated by a correlation coefficient of 0.88.

	Selections as 5-,	8-, and	l <u>13-year-old</u>	Plants	in 1953, 1	956 and 19	961.
Progen y 8	& Survival ¹	Blue S	dlgs ²	Blue	Av	erage hei	ght
Selection	1961	1953	1961	rating	1953	1956	1961
(no.)	(%)	(%)	(%)		(cm)	(cm)	(cm)
42-8	80	55	93	2.7	14	29	60
40 C-1 8	48	3	83	2.1	16	25	67
47-19	- 57	0	80	2.0	14	20	71
40 C- 223	72	9	89	2.2	16	27	81
RC-3	100	25	76	2.2	17	38	84
00-1	96	27	100*	3.3*	14	30	90
40 C- 249	84	2	100*	2.3	17	34	104
PC-4	96	9	83	2.9	16	33	105
40 C-41	96	5	96	2.2	17	37	110
EC-6	91	7	69	2.3	14	38	113
40 C- 24	83	16	62	2.1	16	35	121
EC-1	97	5	97	2.6	13	35	124
42-7	100	10	61	2.1	15	43	128
400-13	97	21	97*	3.6*	15	39	130
40 C 25	100	12	96	2.1	16	39	132
42-6	94	36	73	2.7	17	41	134
40 C-81	100	17	91	2.6	17	38	134
RC-5	100	18	89	3.0	17	37	137
40 C-1 0	91	20	89	3.0	16	40	138
40 C- 22	100	8	93	3.3*	17	42	140
WC-3	97	29	65	2.8	16	42	143
40 C-1	100	4	97	2.3	17	41	149
EC-2	100	10	58	2.6	16	40	150
<u>40C-221</u>	97	19	100*	2.7	20	50	169
Means	91	11	85	2.3	16	37	117

Table 6. Performance of 24 Open-pollinated Progenies of Colorado Spruce

Plant survival as % of original planting in 1954

²Percentage of plants with any evidence of Blue needle coloration ³Average blue rating of plants on a l(green) to 5(intense blue) basis in 1961 *Dominant blue needle-color factors carried by seedtree

It would appear the potential of Colorado spruce progenies could be estimated from the height of the plants at 8 years of age. This may provide a short cut for the early evaluation of the vigor potential of specific hybrid combinations. Two selections (RC-1 and 40C-13) seem to carry dominant factors for blue needle color and produce only blue progeny averaging 3.3 to 3.6 for color density.

<u>Viability of spruce pollen</u> was determined in 1961 on 2% water-agar media with four levels of sucrose content. Fresh pollen from two trees of Colorado spruce and a mixture from several white spruce were dusted on the agar media in petri dishes and incubated at 70°F for 24 hours. Then the dishes were flooded with aceto-carmine stain and the germination recorded with the results listed in Table 7. It was evident the addition of sucrose to the water-agar media increased germination and so provided a more reliable estimate of pollen viability. The results suggest the best germination medium tested for spruce pollen was 2% water-agar with 5% sucrose.

WIGH FO	M. Perera OI DUCIOS	be •			
Sucrose	Colora	ndo	White		
levels	RCL	NC5105	(mixture)	Means	
(%)	(%)	(%)	(%)	(%)	
0.0	23	32	28	28.8	
1.0	29	45	45	39.8	
5.0	53	75	75	69.1	
10.0	51	70	76	65.0	
Means	39	55	56	50.7	

Table 7. Germination for Pollen of Colorado and White Spruce on Water-agar Media with Four Levels of Sucrose.

In another study, pollen from two selections (P-9& NC4-60) was dusted on sucrose-agar media in petri dishes and incubated at 70°F and 90% humidity for 6 to 60 hours. Records for the resulting germination and growth of the pollen have been summarized in Table 8. From these data it is evident that germination of one pollen (NC4-60) started 14 hours earlier than that for the other (P-9). On the other hand, the rate of pollen tube growth for the latter was greatest after 27 hours of incubation, and for the former after 31 hours. No increase in the germination capacity or viability was evident for either pollen after 22 hours of incubation.

Table 8. Viability and Growth of Pollen from Two selections of Colorado Spruce on Sucrose-agar Media after Six to Sixty Hours of Incubation

Hours after	e-agar Meura aro Via	bility ¹	Grand Grand	owth ²
pollination	P-9	NC4-60	P-9	NC4-60
·	(%)	(%)	(u)	(u)
6	Ó	0	0	0
8	0	36.3	0	15
10	0	63.6	0	28
12	0	63.6	· 0	41
22	50	90.9	41	96
27		90.9	88	122
31	60	90.9	122	166
36	60	90.9	149	209*
46	60	90.9	204*	268*
54	60	90.9	219*	287*
60	60	90.9	219*	301*

¹Percentage germination for same field of pollen ²Average length in microns of pollen tube for same field ***Some mold and tubes bursting** A second viability test was conducted using pollen from all 39 seedtrees involved in the 1960 breeding program. Pollen samples were cultured on a sucrose-agar media and incubated for 24 and for 48 hours, after which the pollen was stained with aceto-carmine and the germination recorded. Germination after 24 hours of incubation ranged from 20 to 80% for a mean of 56%, and after 48 hours incubation ranged from 18 to 93% for a mean of 56%. Germination data for 24 and 48 hours were highly associated as indicated by a correlation value of .885. As a result of these two pollen studies it was evident viable pollen was used for all controlled pollinations of Colorado spruce in 1960.

<u>Self-pollination seed</u>, which was produced by 27 self-compatible trees of Colorado spruce in 1960, was sown in the greenhouse in January 1961. Only one of these trees, PC-26 was found to carry the recessive factor for albinoism, and the selfed seed produced 27 green to 8 albino seedlings. Survival and height data were recorded for two inbred and four hybrid progenies of Colorado spruce, which were started in the greenhouse in 1957 and transplanted to the field in 1959. The hybrid progenies of both selections manifested four times the survival and from 20 to 134% greater height growth than the respective inbred progenies.

Efficiency of seven types of isolation bags for spruce breeding were investigated in 1961. Three weaves of cotton and four treatments of cellulose bags were compared for isolation and controlled pollinations on two trees of White spruce. Two of the three cotton bags provided ineffective isolation, while all treatments of cellulose bags proved highly efficient. Cellulose bags, which had been painted with aluminum, proved most satisfactory. This treatment prevented excessive temperatures within the bag, being only 83°F as compared to 92°F for clear cellulose and 81°F for open air temperatures under the extreme conditions prevalent in 1961. Seed set within the aluminum-painted cellulose bag was generally superior to all other cellulose treatments, but on the other hand only approached 50% of that for natural open-pollination. It would appear the southern exposures

of cellulose isolation bags should be painted with aluminum for future spruce breeding work to offset adverse effects of high temperatures from sun radiation.

Germination capacity of Colorado spruce seed at four temperatures was investigated for four seedtrees and four pre-sowing treatments with the results listed in Table 9. It was evident that moisture content of the seed was not materially increased after 15 days of stratification at 41°F even by soaking in water. Both stratification and soaking treatments improved germination capacity of spruce seed, but only under certain temperatures. Generally, stratification proved a superior treatment to soaking for seed of Colorado spruce. Germination for non-stratified seed of Colorado spruce was equal to that for stratification at a temperature range of 75 to 80°F, and inferior at temperatures from 54 to 71°F. Thirty days of stratification is required if Colorado spruce seed is germinated at temperatures below 67°F. Temperatures of 89 to 95°F completely inhibited the germination of this seed.

Four Pre-sov	ving Treatments.			1	
Seed	Seed ¹	Tempe	ratures ²	and Germination ³	
treatments	Moisture	<u>93°</u>	770	68°	61°
	(%)	(%)	(%)	(%)	(%)
nil (check)	6,6	0	91	75	17
Strat. 15 days	31.2	0	91	90	91
Strat. 30 days	32.3	0	95	88	96

78

0

62

74

Table 9. Average Germination for Colorado Spruce Seed at Four Temperatures After

Moisture content of seed after treatment

²Temperature ranged from 89-95, 75-80, 62-71 and 54 to 66°F, respectively ³Germination capacity as % emergence in 35 days

32.6

PROVENANCE TRIALS

Soak 30 days

Two co-operative provenance studies of Scots pine were planted to the field in 1960 and 1961. Some 2,000 seedlings involving eleven provences from M, Holst of Petawawa were planted in 1960 into 20 replications of single-plant plots and three replications of 50 plants per plot, Seedlings of the provenance from Orel, Russia, have exhibited the greatest vigor and survival to date. A

1961 planting involving 25 replicated single-plant plots of 31 Russian provenances was destroyed by drought conditions and replanted in 1962 with an average survival of 92%

Only 1.9% of a 1960 planting, involving 600 seedlings each of 13 <u>Norway spruce provenances</u> from M. Holst, have survived to date. This planting suggests that none of the Russian-Swedish-Siberian-Polish strains of Norway spruce involved can survive the extremes of a mild winter (1960-61) and summer drought (1961) in the prairies even under irrigation.

Nineteen provenances of White spruce from Dr. H. Nienstadt of the Lake States station were transplanted under irrigation in 1960 with an average survival of 17% to date. This material has been soverely ravaged by a mild winter and extreme drought. Results to date suggest some provenances of Quebec, Ontario and New York have greater tolerance to adverse climatic conditions than those of Saskatchewan and Manitoba.

A high <u>mortality of conifers</u> in plantings and natural stands appeared to be general throughout the prairie provinces in 1961. Losses were serious in old established plantings, especially for White spruce and to a lesser degree for Colorado spruce and Scots pine. These results suggest Scots pine may be more resistant to prairie drought conditions (high temperatures, salinity and insects) than are White or Colorado spruce. Research has been undertaken to determine the factors inducing needle-burn and mortality of conifers in this region.

Biennial Report to Committee on Forest Tree Breeding in Canada

May 1962

B. W. Dance Laboratory of Forest Pathology Southern Research Station Maple, Ontario

Testing Poplars for Diseaso Resistance

In 1960 and 1961, techniques were tested for inducing disease in poplar species and varieties, and for assessing the resistance of these hosts to different pathogens whose life histories were studied. These tests were conducted at the Petawawa Forest Experiment Station, Chalk River, Ontario.

Emphasis was on three pathogens: <u>Glocosporium</u> sp., <u>Hypoxylon pruinatum</u> (Kl.) Cke., and <u>Dothichiza populea</u> Sacc. & Briard. However, studies on other potentially import pathogens either were kept active, or initiated.

1. <u>Glocosporium</u> sp. on <u>Populus</u> <u>alba</u> X <u>grandidentata</u>

.

Studies were conducted to determine when <u>Plagiostoma populi</u> Cash & Waterman, the suspected ascigerous state of this foliicolous, imperfect fungus, discharges ascospores suspected of initiating springtime infections. Starting on June 23, 1961, spore traps were set out daily in the presence of perithecia-bearing, overwintered leaves of the hybrid host. This date was selected because, in the previous year, microscopic observations indicated that ascospores did not mature until mid-June.

While the ascospores were trapped regularly in moderate numbers, no peak period of spore dispersal was detected. Instead, with only minor variations, the numbers of ascospores trapped daily declined steadily until July 24-28 when no spores were obtained and trapping was concluded.

Therefore it was considered likely that, contrary to earlier indications, most spores were released prior to June 23, and that infections probably occurred sometime between leaf emergence in May and mid-June, before the majority of leaves matured. Trapping will be conducted in late May and early June, 1962, in an attempt to confirm this conjecture. Meanwhile, data will be collected, as previously, on rainfall, temperature, and relative humidity.

It was determined that sporulating acervuli develop from mid-August onward for an undetermined period prior to leaf fall. However, since no new infections resulted from the micro-conidia released, it was concluded that these spores have only a spermatial function and, therefore, that the fungus requires the entire growing season to complete one cycle of the parasitic phase of its life history.

From these observations it seems evident that ascospores only, rather than conidia, are likely to be effective in starting artificially induced infections désigned to demonstrate variation in disease-resistance between different clones of <u>P. alba</u> X grandidentata.

2. <u>Hypoxylon pruinatum</u>

Artificial inoculations were carried out with this pathogen to determine which types of inocula and infection techniques are effective in starting artificially produced infections on <u>Populus tremuloides</u> Michx., <u>P. grandidentata</u> Michx., and <u>P. tremula</u> L.

Inocula used were ascospores, conidia, and mycelia. Ascospores were discharged directly from perithecia onto exposed cambia; conidial suspensions were injected into unwounded bark, or painted onto exposed cambia and crushed bark; and, mycelia in agar were placed on exposed cambia and covered with moist cotton and tinfoil.

In November, 1961, a series of inoculations was made with mycelia inserted into bark wounds. This test was intended to show if, as has been reported, trees are most susceptible to infection during periods when the bark moisture content is low.

Of 53 inoculations with <u>H. pruinatum</u> made in 1959, 12 may have caused infections by late 1961. In six of these, cankers were well-advanced (1-footlong) and nearly girdling the stems. Furthermore, stromata had formed with perithecia and maturing asci. In every instance, the control remained healthy. These cankers had resulted from mycelial inoculations, and were consecutive in the same series. It seems significant that these most successful inoculations were made in mid-September, and constituted the last of the series that year, because bark moisture content at that time probably was lower than in trees inoculated carlier and which remained healthy.

Recovery of the ascigerous stage from inoculations, and the healthy condition of the controls in late 1961, proves that these artificial inoculations were effective in starting infections. Furthermore, the observed infection period of more than two years agrees with the results of other investigators.

It may be concluded from these tests that ascospores, conidia and mycelia of <u>H</u>. <u>pruinatum</u> are ineffective in starting infections if they are used as inocula on wounded bark during periods of high bark moisture content and with the inoculation techniques used in these experiments. However, it is indicated that a high percentage of infections can be secured with mycelia in wounded bark if the inoculations are made in mid-September, and probably subsequently, when the bark moisture content probably is decreasing.

In 1960, twenty-one inoculations were made with <u>H. pruinatum</u> in the field, on saplings of trembling and large-toothed aspens. Conidial spore suspensions were brushed onto bruised bark. These were examined in August, 1961, but no infections had developed.

In 1961, 70 inoculations were made in the field using conidial and ascosporic suspensions. These were injected into nail holes in the park. By November, incipient infections were evident.

3. Dothichiza populea

en al 👫 👘

A sty it is

To confirm and extend the significant results obtained in 1959 with this pathogen, a massive testing program was executed in 1960. Three hundred and eight, terminal shoots were inoculated in the field with conidial spore

suspensions prepared from pycnidia that had developed on the 1959 series of inoculations.

In 1961, most of these inoculations were microscopically inspected for fruiting structures of the pathogen. With only a few doubtful exceptions, the inoculations were uniformly negative.

No explanation is evident for this reversal of results from those previously obtained. However, defective inoculum may have been responsible.

In the 1961 series, the following poplar species and varieties were inoculated:

Populus tremuloides, grandidentata, balsamifera, deltoides, candicans, tremula, canescens, alba, nigra, Simonii, tristis, koreana, generosa, Rasumowskyana, cathayana, berolinensis, Jackii, robusta, alba X grandidentata, Maine, Geneva, Gelrica, Vernirubens, Carolina, Roxbury, Strathglass, Northwest, Calgary and Brooks.

4. Other Studies

Investigations were continued, or initiated, on fungi having **incom**pletely understood life histories.

(a) Progress was made in establishing the taxonomic status of a new <u>Venturia</u> sp. on <u>P. alba</u> X grandidentata.

(b) A lethal stem canker on large-toothed aspen, caused by <u>Nectria</u> sp? was studied.

(c) The ontogeny and morphology of <u>Marssonina populi</u> (Lib.) Magn., prevalent on <u>P. alba</u> X grandidentata, were investigated.

(d) The orange stain and cambial streaking conditions in clones of <u>P. alba X grandidentata</u> heavily attacked by <u>Gloeosporium</u> sp. was studied.

PHYSIOLOGY OF FLOWERING AND CONE PRODUCTION IN DOUGLAS FIR, 1960-61

by L. F. Ebell

Two replicated, randomized block fertilizer trials using single tree plots and containing 12 treatments had been established in July, 1957, partially refertilized in May 1958, and all trees refertilized in May 1959, in flower induction experiments. The July, 1957 applications were too late for flower bud formation, and although the effective treatments of the partial 1958 refertilization produced double the number of cones as the non-refertilized portion, no statistically significant cone induction response was obtained in 1959, a near record cone crop year. The nitrate nitrogen treatments of the 1959 refertilization produced highly significant cone crop responses in 1960 despite absence of natural cone production. Ammonia nitrogen treatments did not significantly elevate cone production until 1961, the second season after refertilization. Since growth responses were the same for both forms, it seems probable that nitrate nitrogen may promote specific metabolic responses favouring floral initiation. In 1962, the third season after refertilization, both forms produced highly significant cone crop responses, possibly indicating an increased turnover and elevation of nitrogen in the nitrogen cycle. Significant regressions of cone count on diameter growth of the previous season were obtained for the 1959-1961 cone crop years, but while the best cone production occurred on trees with average or better-than-average growth, the fastest growing individuals produced few or no cones. The basic data has been obtained for further investigation of this regression on material not confounded by cone induction treatments. A report, for publication, on the results to date from the fertilizer experiments has been prepared.

Biochemical analysis, conducted to date, on branch tissues from check and girdled trunks of double-trunked Douglas fir indicate heavy cone crop production induced on the girdled trunk was not preceded by higher levels of free sugars but was preceded by elevated starch food reserves. Part of the photosynthates confined to the crown by girdling were diverted to increased diameter growth. A specific micromethod of starch analysis, which is not influenced by hemicellulose pentosans or variations in amylose : amylopectin ratios due to anatomical or seasonal factors, has been developed for this work.

Two of six Douglas fir clones established by grafting in 1959 contained male and female cone-producing ramets in the spring of 1962. One of these clones was from an exceptionally heavy and reliable cone-producing parent tree, the other from a good cone-producing parent at the elevational limit of Douglas fir on a central Vancouver Island mountain. No evidence of flowering or cone production was found on the clones representing barren parental material.

Some members may recall that the first evidence of rooting of June, 1959 air-layers was uncovered at the August 1960 meeting at Lake Cowichan. Further examination revealed a rooting success of 20-40% on three trees on which layers had been placed on 2-3 year old branch wood. The layers on two other trees suffered from extensive breakage and suppression due to placement on thinner, one-year-old branch tissue. Although no further trials have been made it would seem essential to place Douglas fir air-layers no later than the latter part of April in order to obtain rooting within a single season. Due to extremely rapid callusing and bridging in the early spring it might then be necessary to re-girdle and re-apply rooting hormones. Pollen sampling studies conducted in the 1958-1960 period have been concluded with the report cited below. One important conclusion is that it is unlikely that rainy weather can seriously impede pollination success. The bulk of the pollen of all species was invariably shed in only four to five days interspersed throughout the flowering period, regardless of whether the season was abnormally wet or unseasonably dry.

Submitted for Publication:

Ebell, L. F. and R. L. Schmidt. Influence of meteorological factors upon coniferous pollen production and dispersal on Vancouver Island, British Columbia.

APPENDIX G

Report to Committee on Forest Tree Breeding in Canada, July 1960-June 1962.

Faculty of Forestry, University of Toronto

by J. L. Farrar

1. Introgressive Hybridization in Picea rubens and P. mariana.

This study began with a comparison of twig characters of the progeny of certain stands grown by Mr. Mark Holst at the Petawawa Forest Experiment Station, Canada Department of Forestry. It was extended to include a field investigation of certain of the parental stands and other trees located from North Carolina to the University Forest in Haliburton County, Ontario. The study indicated that consistent differences exist between the two species with respect to morphology of fruit and twigs, and ecological relations. No differences were found in chromosome morphology. Hindrances to hybridization were attributed mainly to ecological isolation. The two species have likely co-existed since the Pliocene epoch. The variation between various populations was analyzed by Anderson's hybrid index method and pictorialized scatter diagrams. The analysis indicated that introgressive hybridization has occurred between the two species. In Quebec, black spruce seems to be essentially pure with the direction of gene flow being toward red spruce. In the Maritime provenances gene flow has occurred in both directions with a somewhat greater flow toward black spruce. Little evidence of introgression was found in the Northeastern region of the United States. The study is the subject matter of the thesis submitted by Mr. Morgenstern as one of the requirements for the Master's Degree in Forestry.

2. A Provenance study of seedlings of Pinus banksiana

Seedlings of nine provenances of jack pine from across Canada were germinated and grown in a growth chamber. The more northern provenances were more responsive to photoperiodic treatments than were the southern ones. The total dry weight, height, dry weights of leaves and roots, and dry weight per unit of nitrogen content were all positively correlated with the number of growing degree-days (base 42° F) in the locality of origin. The study was carried out by Mr. M. M. Giertych, using seed lots supplied by Mr. Mark Holst.

Publications

Giertych, M. M. and J. L. Farrar. 1961. The effect of photoperiod and nitrogen in the growth and development of seedlings of jack pine. Can. Jour. Botany 39:1247-1254.

BIENNIAL REPORT - HARD PINE BREEDING

1960 and 1961

D. P. Fowler

Ontario Department of Lands and Forests, Research Branch, MAPLE, Ontario.

The development of new types, resistant to the European pine shoot moth and superior in growth form and growth rate continued to be the aim of this project. During the academic year 1959-60 the writer was on educational leave from the Department of Lands and Forests. During this period the hard pine work maintained its status quo. No hard pine pollinations were made in the spring of 1960. In 1961 the pollination program was almost completely devoted to a study of the effects of inbreeding in red pine, <u>Pinus</u> resinosa.

<u>Acquisitions</u>: During 1960 and 1961 the following new clones and populations were acquired:

	<u>Species</u>	No. clones	No. populations	
Ρ.	resinosa	25	20	
P.	leucodermis	2		
Ρ.	radiata	5		
Ρ.	nigra austriaca x silvestris	3		
P.	rigida x taeda		1	
		<u>35</u>	<u>21</u>	

<u>Seeds and seedlings from 1959 pollinations</u>: Controlled pollinations carried out in 1959 produced the following seeds and seedlings in 1960 and 1961 respectively:

and fight the later of the second states of the second second second second second second second second second

Bo Yere W. C. C. Star Die LAND A MEL STOLE Charles de Listado

 $g_{1} = -\infty e_{1} g_{1} + \cdots + g_{n} = -\infty e_{n} g_{1} + \cdots + g_{n} = -\infty e_{n} g_{1} + \cdots + g_{n} = -\infty e_{n} g_{n} + \cdots + g_{n} = -\infty e_{n} g_{n}$ $\sim p_{\rm eff} \sim 1000$ and ~ 1000 enter and the second ender is proved in the constant of the firm () and the light of the light of the second state of the the construction of the second beaution of the second second second second second second second second second s where $g_{0} = g_{0} = g_{0}$ (0,0) (0,0) (0,0) (0,0) (0,0) (0,0)

georgian and the second s an an tha tha Ministra There are a start of the second second

وه ۲۰۰۰ ۲۰۰۰ ۲۰۰۰ ۲۰۰۰ میرمد و میرو

(a) A second s second s Second sec teste e E 🖓 en avere en el composition de la composition de er en en en en art sollt
Cross (No. Crosses	No. Flow- ers	No. Cones	No. Full Seed 1960	No. Seed to Germ	No. Seed- lings . <u>1961</u>	
<pre>(densiflora x austriaca) x resinosa " x silvestris " x self (austriaca x densiflora) x self resinosa x (austriaca x densiflora) silvestris x (densiflora x austriaca) silvestris x (austriaca x densiflora) nigra x silvestris silvestris x silvestris (early flowering) rigida x taeda</pre>	1 1 1 1 5 1 1 2 1	118 98 99 98 79 111 25 51 103 27	52 51 44 84 41 84 18 28 	36 577 28 77 10 576 3 12 732 43	36 100 28 77 10 362 3 12 200 43	32 98 24 71 7 317 317 3 10 105 43	· · · ·

Effect of inbreeding: In the spring of 1961, 150 different red pine pollinations were made on 2068 flowers at Kane, Pennsylvania, the Lake Abitibi area, and Swastika and Chalk River, Ontario. The following pollinations were made:

Location	Origin	No. Trees		Num 	ber flo <u>Typ</u> Self	owers <u>e Pol</u> Mixed Local	pollina <u>len</u> Mixed Maine	ted	
Kane, Pa.	East. U.S.	4			55	56	57	· · · · · · · · · · · · · · · · · · ·	
11	N. W. Minn.	4			58	58	62		
9 3	Central Wisc.	4			64	56	61		
89	N. E. Minn.	4			44	56	51		
" N.E.Wis	cS. Upper Mich.	4			43	47	47		
11	Head of Lakes	4			48	46	47		
9 9	Brainerd Cameron	4			39	46	40		
3 6	N. Upper Mich.	4			43	47	45		
5 8	Lower Mich.	4			56	53	52		
Northern Ont.	Lake Abitibi	4			59	58	53		-
Central Ont.	Chalk River	5			78	111	89		
Northern Ont.	Swastika	5			6 6	86	91		
		TOTAL			653	720	695	2068	
Central Ont.	Algonquin Park	2	76	flowers	bagged pollir	l but nated.	not		

	.01 86.58 11086 1261	つ わた」考 (13) (注)。		.0) •	• 12 ¹ • 89 N_ 11 14 • 3	: * • ; *	
••••	يعودت المتحدوث كالم		× 4 · 2 · ·			. Se adama a	and a second
			•	200		P	and the second
		1.5					
	$\sum_{i=1}^{n}$			33 ³	1. A. A.		and the second sec
			<u>,</u> ,				A CONTRACTOR OF MELANDER CONTRACTOR
			1.5	ia.			,这些你们的是我们的。""你们的你们的你们,你们就是你说。"
	· · ·	÷.,	· ·				1991年1月1日日前的1月1日日的1月1日日(1991年1月1日日) 1991年1月1日日(1991年1月1日日)(1991年1月1日日)
	:					i	rendirende al de l'ante i d'arte e e
		5 8					化合金合金 化合金合金合金合金合金合金合金合金合金合金合金合金合金合金合金合金合
							and the second
	691		1. 1 N		· · ·		q = 2 + 1 , $3 = 2 + 2 + 2$, $q = 1$
			14 - 44	۰.	: `		$\sum_{i=1}^{n-1} (1-i) \sum_{i=1}^{n-1} (1-i) \sum_{i$
		· · · ·					

	- 3	n Frida Santa Santa Frida Santa	ing a start of the	ad de		
ò.`		- , ' *_				
		· .				
						general terms of the second
						and the second
	Ċ.					$\{\phi_{i},\phi_{j},\phi_{i},\phi_{j},\phi_{$
						1. (1. (1. (1. (1. (1. (1. (1. (1. (1. (
					,	
		1	~			n an an ann an ann an ann an ann an ann an a
		· · · *			÷.,	
						station of the second sec
	· .					 A state of the sta
	-					

First cycle measurements were made on seedlings resulting from self and cross pollinations on five red pine trees at Vivian, Ontario in 1958. The results indicate that selfing had little or no effect on flower, cone, seed or early growth characters. These five trees are self-fertile and self-compatible. Seedlings resulting from self and cross pollinations of four additional trees pollinated in 1959 were not measured in 1961. Flower, cone and seed characters were not influenced by the type of pollen used in these crosses.

A small preliminary study of the effects of inbreeding in jack pine, <u>Pinus banksiana</u>, was initiated. The following crosses were made at the Southern Research Station:

<u>No. trees</u>	No. flowers	pollinated
	Self pollination	Cross pollination
5	136	128

<u>Selective fertilization in Pinus resinosa</u>: Two trees, one (#585), a known carrier of a recessive marker gene, and the other (#622), a possible carrier of this same gene, were used in this work. The following crosses were made in 1961:

Cross	No. flowers pollinated
585 x 585 (self)	9
mixtures tested)	35
622 x 622 (self)	24
622 x 585	2/

<u>Vegetative propagation</u>: A grafting experiment, to determine if the cold period requirements of red pine could be satisfied by storing fall collected scions in a deep freeze, was carried out. Red pine scions were collected in early November of 1960 and stored in a deep freeze at -20°C. Stored and freshly collected scions were grafted on scots pine stock at two-week intervals from November 1960 to February 1961.

No significant differences were found (anal. of variance and anal. of covariance) between grafts made with stored and fresh scions for either survival or shoot production. Analysis of covariance, treating the relationship between survival, number of shoots and date as linear regressions, showed a significant regression between survival percentage and date (F = 4.90) and a highly significant regression between percentage of grafts producing shoots and date (F = 14.73). Survival percentages were 78% before January 1st and 46% after, and percentage producing shoots 57% and 12% respectively. It would appear from the experiment that the earlier grafting is done in the fall, the better, and that storage of scions in a deep freeze does not improve subsequent survival and growth.

Turkey Point planting area: The following clones and populations were planted at Turkey Point, Ontario in 1960 and 1961:

Species	No. Clones	No. Plants	No. Pops	No. . Plants	Purpose
P. resinosa	8	104			Plus trees.N.Ont.
			1	17	Observation
			6	270	Half-sib progeny
					test
P. nigra			5	360	Provenance test
			14	1172	Observation
P. silvestris			13	888	Freedom from
					Cronartium
	_		8	1750	Observation
P. thunbergii	2	17	2	876	Observation
P. densiflora erecta			1	583	Observation
P. densitlora x silvestris			10	389	Observation
P. silvestris x (densiflora	x austr	iaca)	2	107	Observation
P. densiflora x thunbergii P. nigra cebennensis x			1	440	Observation
densiflora	1	1	1	4	Observation

H-4

Report to the Committee on Forest Tree Breeding in Canada

Tree Physiology Section

Petawawa Forest Experiment Station

D.A. Fraser June 1, 1962

Apical and radial growth, and the developmental anatomy of vegetative and reproductive buds of white spruce were investigated. Apical growth of the leader occurred from mid-May until mid-July, that of the side branches from mid-May until late June or early July. Total apical growth of the side branches, in general, diminished with their distance from the leader. Initiation of radial growth started in the main axis behind the terminal bud and moved basipetally along the trunk. Cessation of radial growth followed the same order.

The apex of the vegetative bud which developed in late July or early August includes a central zone, a peripheral zone with needle primordia, and a basal zone of thick-walled cells. Reproductive buds could be recognized by early August as oblong structures with globose appendages. By mid-August, these appendages had differentiated into either pollen sacs with pollen mother cells in the male buds, or into bracts and ovuliferous scales in the female ones. The female cone entered winter with little visible internal differentiation. The formation of reproductive buds in white spruce was associated with higher than average day and night temperatures, together with more than average sunshine (usually associated with less rainfall).

Black and white spruce seedlings were grown for two years in growth chambers under extended photoperiods. Under continuous incandescent light, the black spruce grew 30 inches whereas the white spruce grew only 5 inches. Growth of both species was somewhat greater in the greenhouse, where the continuous light consisted of natural daylight plu light from a combination of fluorescent tubes and incandescent bulbs during the night period. Black spruce seedlings grown under continuous light in the laboratory during the winter were successfully transferred to an outside nursery in the spring and assumed there the normal seasonal periodicity of growth. Black spruce plants that were grown completely under a 24-hour photoperiod formed a terminal bud when transferred to a 16-hour photoperiod. The dormancy of this bud was broken when the needles were killed by high temperatures; apical growth continued then until new needles were formed. Apical growth of black, white, and red spruce seedlings in the nursery was significantly increased by a 24-hour photoperiod effected through supplementary illumination during the summer months.

2

Growth regulators (promoters and inhibitors) are being investigated by paper chromatography and <u>Avena</u> coleoptile assays. There was a high content of growth promoting substances in actively growing terminal buds with lesser amounts in lateral ones. The amount decreased as apical growth ceased in July and the terminal buds were formed. A growth inhibitor was extracted from dormant buds. This inhibitor also appeared in spruce and birch seedlings, when they were moved from long to short photoperiods in the growth chamber.

Paper chromatographic analysis of the ethanol extracts of spruce buds continues to elucidate the quality and quantity of free amino acids at different stages in the development of the apical meristem.

Oxygen uptake by buds of white and black spruce was determined in a microrespirometer. Increase of respiratory activity in the terminal buds at the top of the tree in May coincided with initiation of apical growth. The respiration of shoots destined to produce either vegetative or reproductive buds will be studied this summer.

Biochemical assay of bud homogenates was conducted to determine presence of enzymes considered active as terminal oxidases. Cytochrome oxidase activity . was low. Ascorbic acid oxidase and polyphenol oxidase were present as active enzyme systems.

Publications:

FRASER, D.A. 1961. Transport of mineral nutrients in trees. <u>In</u> Recent Advances in Botany, 1311-1316, University of Toronto Press.

> 1961. Plant science in the service of man. The Crucible (Easter), 10-12. Abstract from a lecture for the annual meeting, Ontario Secondary School Teachers Federation, Dist. 25, Richmond, Ontario, October 28, 1960.

_____1961. Tree growth in relation to soil moisture. <u>In</u> Tree Growth, <u>Edited</u> by T.T. Kozlowski, The Ronald Press, New York, 183-204.

1962. Growth of spruce seedlings under long photoperiods. Canada Dept. of Forestry, Forest Research Branch, Tech. Note No. 114, 18pp.

1962. Apical and radial growth of white spruce (<u>Picea glauca</u> (Moench) Voss) at Chalk River, Ontario, Canada. Can. J. Botany 40: 659-668.

and GAERTNER, E.E. 1961. Microenvironmental studies in Canadian forests. <u>In</u> Recent Advances in Botany, 1391-1396, University of Toronto Press.

and GAERTNER, E.E. 1961. Use of radioisotopes in forestry research. In Recent Advances in Botany, 1381-1387, University of Toronto Press.

REPORT TO COMMITTEE ON FOREST TREE BREEDING

and the second July, 1962

W.V. Hancock

Department of Forestry of Canada, Forest Products Research Branch, Vancouver Laboratory

. . The Assessment of Wood Quality of Trees Selected for Breeding Purposes

The program, as described in the report presented at the Seventh Meeting of the Committee for Forest Tree Breeding held at Cowichan Lake, B.C., August 21-23, 1960, has been continued with some changes in technique and a reduction in intensity of sampling and measurement.

Tracheid Length

Tracheid lengths were measured in detail in thirty-one trees. The normal pattern was a rapidly increasing tracheid length with an age from the pith followed by a period of less rapid increase. Based on age only without regard to growth rate, trees with high initial tracheid length did not always have longer tracheids at some distance from the pith than trees with shorter initial tracheids. Further study of the data is needed to clarify these relationships for the trees under study.

Because all of the trees studied rapidly reached a length of tracheid considered suitable for pulp and because of the great amount of routine work involved, further tracheid length studies will be confined to those trees for which progeny are available and on which the determinations of inheritability will be based.

Spiral Grain

Spiral grain was measured from pith to bark in 177 trees that are potential breeding material or are being used for genetic studies. It was not measured in all of the associated dominant trees collected along with the potential superior or plus trees. These associated trees were examined for purposes of comparison with plus trees to determine relative efficiency of site utilization, and comparisons of severity of spiral are of no value.

Within those trees measured an upper limit of 6° of spiral was set which corresponds to the permissible slope of grain of one inch in ten inches set out in lumber grading rules for construction grades. None of the trees selected as potential seed-orchard trees exceeded allowable limits. It was found that earlier measurements based on 0.16 inch diameter cores were frequently in error, apparently due to twisting of the cores during extraction. Cores collected with 12 mm. or 0,75 inch diameter borers appear to give a true read-ing. . '

Compression Wood

The presence of this defect was determined by the light-box method, and was found to be of limited occurrence and generally confined to the first few rings in the plus trees measured. These trees were selected for good form and straightness of stem and compression wood should not occur in them in large amounts. Material containing compression wood was excluded from calculations.

Age. Rate of Growth and Specific Gravity

Initially the age, rate of growth and specific gravity were determined for each radius of diametral core from pith to bark. The first ten years were measured in annual increments and the balance of each oore in five-year increments. A pattern developed that showed great variability over the first ten years, frequently with very high initial values, but falling off to about ten years and then rapidly rising again. The period from ten years on gave relatively regular curves of specific gravity over age with a close approach to the final high level at about thirty-five years, although specific gravity of many trees was still increasing. Consequently the period ten to thirtyfive years was chosen as a basis for comparison. Data from this period were used to draw regression lines for each radius. These were averaged to yield a single regression and, from this, the mid-point (22.5 years) was designated as the point of comparison between different trees. All comparisons were made within the limits of the population which was chosen as including those trees which had been measured. Through further mathematical treatment it was found possible to assign a comparative mathematical value on the scale 1 to 10 to each tree. It is proposed to try to integrate this into a numerical value for all tree characteristics.

Future Work

Recently the work of detailed measurement of collected cores was taken over by the Faculty of Forestry, University of British Columbia, under a grant from the Canadian Pulp and Paper Research Institute. The University is, in addition, undertaking to develop alternative or simplified techniques to speed up the obtaining of information. An analysis of the work done to date in this laboratory is being made.

Inheritability Studies

While the short-term objective of this study was the assessment of potentially superior trees for inclusion in seed orchards, the long-term aim is to study the inheritability of anatomical characteristics of various species. This work is still going forward.

and a second process

. . . .

The inheritability of wood quality is being investigated using several second-growth Douglas fir trees and their progeny from control-pollinated seed, including several apparently apomictic trees. A detailed analysis of the first parent has now been completed. This is a 22-year old tree 11.4 inches d.b.h.i.b. Two problems were studied:

1. The determination of stem tracheid length and specific gravity patterns with regard to age from the pith and height, and

2. The correlation of these features with branchwood.

The ultimate aim of such correlations is the accurate prediction of stemwood quality from branchwood in order to make possible the non-destructive evaluation of plus-tree progeny. Tracheid length and specific gravity were determined for stemwood at five levels, 4.5, 14.5, 24.5, 34.5 and 44.5 feet and for branchwood at the three upper levels. Measurements were recorded individually for the first 10 annual rings and thereafter in groups of five or less. Wood near the pith was measured in greater detail because the first material available from progeny will come from this area.

In general, analyses of variance for specific gravity regressions revealed no significant correlations between stemwood and branchwood, while those for tracheid length showed highly significant correlations. Progeny from this tree are available but measurements have not yet been made.

SUMMARY REPORT ON FOREST TREE BREEDING

1960 and 1961

C. Heimburger

Ontario Department of Lands and Forests, Research Branch, MAPLE, Ontario.

Breeding work with white pine, aspen poplars, hard pines and white cedar was continued. Work with chestnut was reduced to a maintenance basis in 1961.

White Pine

Resistance to blister rust and weevil, and satisfactory growth form and growth rate were the main objectives in breeding. Acquisitions of new materials included scions of Idaho white pine showing high heritability of blister rust resistance, scions of P. albicaulis from Idaho, Alberta, California and Oregon, several P. cembra materials from Switzerland and a few clones of P. flexilis, P. monticola, P. koraiensis and P. strobus. The plantation of P. koraiensis at Orono, Ont. had an abundant cone crop in the fall of 1960 yielding 12 progenies in 1961. Selections from seedling populations screened for resistance to blister rust comprised 59 clones; of these, 30 clones were P. griffithii from high elevation in Pakistan and 25 clones were natural P. griffithii x strobus hybrids from the Arnold Arboretum. Hybridization in 1960 was centered on interracial work with native white pine, using pollen from one good tree at Pointe Platon, P.Q. on grafts of the same origin at Maple. Pollen of a promising

(1) A state of the state of

en and A .

A state of the sta

is the first state

A second seco

and a second and subscriptions of a construction of the second and a start of the s The start of the start energi kara la filo de la construcción de la const and the second energy standy affiliate and the standard standard and the standard standard standard standard standard standard Condition of the state of th THE REPORTED BY ALL REPORTED AND REPORTED BY AND REPORTED AND A REPORT OF A REPORT OF A REPORT OF A REPORT OF A Next of the second of the second s the second second provide a second for the second second as an and the state provide the second provided the graves grafie and a the set of the later of the set of the set

P. griffithii was received from Rochester, N.Y. and much of our pollen was used to determine the breeding value of our materials in respect to resistance to blister rust. A total of 86 intraspecific and interspecific crosses made in 1960 yielded 4,033 full seeds. A fairly large seed collection from resistant trees in Germany yielded numerous seedlings that, thus far, do not seem to be more resistant to blister rust than the average native white pine in our tests. The plantation of white pine selected for resistance to blister rust and established at Connaught Ranges near Ottawa, in cooperation with the Department of Forestry, received a heavy thinning in 1960, removing all trees with blister rust attack either directly or on their grafts at Maple. This caused abundant flowering in 1961 and gives promise of a good cone crop in 1962. Three kinds of interspecific crosses were made in 1961, namely 6 crosses of P. monticola x griffithii, 2 crosses of P. armandi x albicaulis and 30 crosses of P. koralensis x albicaulis. The range-wide 5-station provenance test of white pine initiated in 1957 was outplanted in 1961. One outplanting was established at Turkey Point, Ont. with a total of 6,292 trees. Another outplanting, with the same 13 provenances, was established in the Ganaraska Forest and comprised 4,212 trees. The selection for resistance to weeviling, in co-operation with the Forest Insect Laboratory in Sault Ste. Marie, was continued. A plantation of white pine in the Kirkwood Management Unit was top-grafted in 1961 for this purpose with 1050 scions of 21 clones of various native

and hybrid white pine materials selected for resistance to blister rust. An experiment in the propagation of juvenile white pine by stem cuttings, with the aid of growth hormones, was started in the fall of 1960. Next year it yielded 55% rooted cuttings after the application of Rootone, as compared with 3.3% rooting of untreated controls. Other hormone treatments yielded intermediate results. The rooting obtained is too low for routine propagation but is encouraging as a basis for further tests. Experiments in bench grafting were carried out during both winters of 1960-62 and established a good basis for the application of this method to routine propagation in the future.

Some white pine seeds were irradiated with thermal neutrons at the plant of Atomic Energy of Canada Ltd. in the fall of 1960. Two hours in the pile with a total dose of about 6000 rem reduced germination to 36% of untreated controls, i.e. the L 50 dose is somewhat lower than this treatment. Three hours in the pile with a total dose of about 9000 rem reduced germination to 9% of untreated controls and 4 hours to 2%. The surviving seedlings have thus far not shown abnormal growth peculiarities. Their resistance to blister rust in comparison with seedlings raised from untreated seeds of the same origin is being studied. Seeds from several plus trees at the Petawawa F.E.S. were obtained in 1960 and have yielded seedlings for progeny tests, mainly in respect to resistance to blister rust.

Poplars

The production of aspen-like hybrids, suitable for growing

in southern Ontario continued to be the chief aim of this project. Acquisition of new materials was in 1960 restricted to a few clones of Populus alba, P. tremula and P. canescens from Switzerland and In 1961, 10 clones of P. euramericana were acquired Yugoslavia. from central Europe, 8 clones of P. tremuloides from Idaho, British Columbia, New Brunswick and Ontario, and 14 clones of P. canescens from the Netherlands and Germany. Hybridization was in both years centered on the production of second-generation hybrids between \underline{P} . alba and aspens, using several first-generation hybrids and \underline{P} . canescens materials in our collection that are beginning to flower. The breeding and selection of precocious aspen types was continued on a small scale. A good method for testing the rooting ability of stem cuttings was developed and applied to numerous new aspen Several promising clones of silver poplar and aspen materials. hybrids raised at this Station were selected on the basis of this A cross of one of our P. alba with a P. alba from Spain new test. yielded some materials with better rooting ability from stem cuttings than either of their parents, indicating the action of complementary genes in this respect. It was found that crossing of silver poplar of superior rooting ability with aspens varies considerably in the effect on the rooting ability of the hybrids, depending also on the aspen species used. P. grandidentata is one of the best for this purpose. Continued heavy attacks by a fungus believed to be a <u>Gnomonia</u> species is a serious problem in aspen breeding. Asiatic aspens and their hybrids seem to be particularly susceptible to attacks. Hybrids of P.trichocarpa

and a sector model for here that to see that an a sector sector many sector many sector as and the set of the transmission of any state of the set set of the the formation in the second and a second provide the second of the second s t^{ab} , we can also also also be the set of the set There is a press of the closer of the second for the second for the the control of the second control of the second second second to the second second second second second second The memory of the second s The general second second stands and a second se Detroit des de la contra de la contra de contra en contra ante de la contra de la contra de la contra de la co and the second The second se 化酸磷酸钙 化结合化物合物 化放射性放射 法外外的保持上述 建铁橡胶橡胶 法不能分析 化分子子 化水平分子用分子 计操作 生产性的现在分词 化水子工具 化分子性化 计分子分词 化合理合金 化分子分析 化分析化分析 nten and style of the second of the second contract of the second s and the main of the state processor courses when it has a course when the shout the shirt of our grades into a structure and the second second second The second second and a second s . We can be the end of $\sqrt{2}$, the density 2 , the density 2 , the end 2 , patient for which the providence of a second constraint of the department of the and the term the second second second second and the second second second second second second second second se and the second and the second and the first sector of the standard of the sector of the All manifest and the state of a manifest of the state of the

with various forms of <u>P</u>. <u>alba</u> are beginning to show promising possibilities of introducing the superior rooting ability of <u>P</u>. <u>trichocarpa</u> into aspen hybrids. One large test plantation was established in the spring and two smaller ones in the fall of 1960, and two in the fall of 1961.

Hard Pines

See report by D. P. Fowler; H-1 to H-4 White Cedar

The production of improved types of this species, Thuja occidentalis, and of western red cedar, Thuja plicata, and hybrids, hardy in southern Ontario, is the aim of this project. The large collection of western red cedar seedlings was so covered with snow in the winter of 1959-60 that it was not possible to subject them to a good freezing test. Instead, the seedlings were heavily culled in the beds, in respect to size and possible signs of winter damage. It was found that the hardy western red cedar, received previously from Rochester, N.Y. are most probably hybrids with Thuja standishii from Japan, which probably accounts, to some extent, for their hardiness under our conditions. A provenance test plantation, consisting of 15 provenances of white cedar from Ontario, Quebec, New Brunswick, Manitoba and Wisconsin, was established in the Ganaraska Forest in the spring of 1961. In the same year, two observation areas of western red cedar, one of 33 populations and another of 8 populations, were also established in the Ganaraska Forest, in addition to one small observation area of white cedar of 10 populations. A number of rooted cuttings and

hybrid seedlings were set out on their permanent location at this Station to serve as future breeding materials.

Chestnut

The aim of this project is the production of hardy dwarf types, resistant to blight and suitable as dwarfing stock in the breeding program with timber-type chestnuts. In 1960, seeds of Chinese chestnut, from Beltsville, Md. produced plants of good growth and hardiness. The next year these plants were moved from the nursery to various places at this Station for further observation and breeding. Chinkapin seeds were again received from Virginia, but did not germinate. Seeds of European chestnut, obtained in a grocery store in Toronto, yielded seedlings that were not hardy. No new work was undertaken on this project in 1961.

Publications

Fowler, D.P. 1961. The effect of photo period on white pine seedling growth. The Forestry Chronicle 37 (2): 133-143.

Heimburger, C. 1960. Review of "Principles of Plant Breeding" by R. W. Allard, 1960. The Forestry Chronicle 37(1): 67-68.

Heimburger, C. 1961. Comments: Problems in Forest Tree Breeding. Recent Advances in Botany: 1699-1703.

BIENNIAL REPORT

APRIL 1, 1960 TO MARCH 31, 1962

FOREST TREE BREEDING AND GENETICS

AT THE

PETAWAWA FOREST EXPERIMENT STATION

M. J. HOLST

INTRODUCTION

This report gives a brief outline of the work done by the tree breeding group at Petawawa Forest Experiment Station.

Our early efforts start to bear fruit and many experiments have been established in Canada, the United States and overseas. We are still in the "provenance stage," as Canada is pitifully short of adequate provenance experiments. Both foresters and tree breeders will find our provenance work useful but we are looking forward to the day when we can do more breeding work.

WHITE SPRUCE

An early provenance experiment (Exp. No. 93), including 32 races of white spruce mainly from the middle part of the Great Lakes-St. Lawrence Forest Region, has finally been field planted in 21 locations in Canada and the United States. This experiment is a forerunner to the more comprehensive white spruce provenance experiment (Exp. No. 194) conducted in co-operation with the Canadian Pulp and Paper Association, Ontario and Quebec Departments of Lands and Forests, and the University of Toronto. This last experiment is progressing on schedule. Two-year-old stock was distributed for the five western experiments in 1961, and seedlings for the five eastern experiments were shipped in 1962. In 1963 seedlings for four complete experiments will be transplanted in the nurseries at Acadia, Valcartier and Petawawa Forest Experiment Stations, and in the Fort William Nursery (Ontario Department of Lands and Forests). The first field planting will be in 1963.

The pilot white spruce provenance experiment to Exp. No. 93 has been rated annually. Furthermore a study was made of needle and wood dry weights to see whether some provenances have a more efficient photosynthesis than others. This certainly seems to be the case, but no clear pattern emerged. The wood/ needle weight ratios show random distribution.

A number of inbred lines of white spruce were sown and followed closely for mortality, and for albino and yellow seedlings. We hope to be able to study the early occurrence of freak seedlings in relation to later growth. Perhaps we may also be able to estimate breeding value of plus trees by comparing the relative performance of the selfed progenies, and get rid of deleterious genes as we establish inbred lines for future outcrossing. We have also started some grafting experiments with young, middleaged and old white spruces to study the influence of scion topophysis, graft position, and root-stock pruning on the development of the grafts. (Exp.No.241)

Seed was distributed earlier to test Danish Sitka x white spruce hybrids against a number of white spruce and Norway spruce provenances for growth rate and resistance both to the Sitka spruce weevil and the white pine weevil. This material has been planted at the Green Timber Nursery in British Columbia (Exp. No. 129) and in six spruce weevil exposure plots in coastal Washington and Oregon (Exp.No. 127) by J.W. Duffield, Industrial Forestry Association, Nisqually, Washington. He writes that St. Zenon, P.Q. white spruce has better than average growth rate and some of them produce phenomenally . . . straight leaders. This is a surprise as the St. Zenon lot in our experiments has been 14 percent inferior to local white spruce. One experiment was planted at Petawawa (Exp.No. 139) with certain misgivings about its hardiness. However, nearly all of the various Sitka x white spruce hybrids showed pronounced magnesium deficiency - perhaps a good indication to keep these hybrids in coastal In Quebec this material has been planted in five localities (Exp.No. areas. 135); in several localities in New Brunswick and Nova Scotia (Exp.No. 134), and in one locality in New York (Exp.No. 138) and Newfoundland (Exp.No. 136). This experiment should eventually give interesting information regarding performance in a western oceanic, a continental and in an eastern oceanic climate. If the hybrids on the west coast are resistant to the Sitka spruce weevil, this could be a very useful discovery as hardly any Sitka spruce is planted because of the weevil menace. For the east coast we can hope for a fast growing or at least a salt tolerant hybrid.

RED AND BLACK SPRUCE

We have at the Station material indicating that red spruce crossed with Petawawa black spruce produces a heterotic hybrid. This is a valuable lead, but the material originates from the area where the ranges of red spruce and black spruce overlap and where introgression is common. Chances for getting even more heterosis should be good if we use material of both species from outside their overlapping zone. We should also try to balance the climate because the hybrid will most likely be intermediate in climatic preference. If we want to use southern Appalachian red spruce, we should cross it with boreal or western black spruce to make a hybrid suitable for the Great Lakes-St. Lawrence Region or the Acadian Forest Region. This is the idea behind our hybridization work in red and black spruce.

The species flowered abundantly in 1960 and we concentrated on making black and red spruce hybrids. To that end Messrs. C.W. Yeatman and K. Morgenstern went to the United States and collected large amounts of red spruce pollen from Pennsylvania, West Virginia and North Carolina. This work was only partly successful because most of the pollen came too late to be of any use at the Petawawa Forest Experiment Station and the crosses were therefore repeated in northern Ontario at Cochrane.

In 1961 there were hardly any black spruce flowers in Ontario. By looking at climatic maps, we located an area in Manitoba that had above average spring temperatures in 1960, and a phone call to our District Office in Winnipeg confirmed that black spruce was flowering abundantly. So we repeated the red spruce crosses on black spruce in Sandilands Forest Reserve and Riding Mountain National Park.

All in all this pollination work has been a disappointment. Although the red spruce pollen germinated well (from 35 to 65 percent), we got very small cones and extremely low germination. This may be due to the following:

(1) the pollen germination test is a poor indicator of pollen viability; or

(2) our crown isolation tents are unsuitable for black spruce; or
 (3) black spruce outside the overlapping zone is incompatible with r

3) black spruce outside the overlapping zone is incompatible with red spruce.

To solve these problers we repeated these crosses on Petawawa black spruce with old and fresh pollen in 1962.

Our red spruce provenance experiment with 22 races (Exp.No. 95) has now been field planted in 23 locations in Canada and the United States. Seed of some of the provenances has been distributed overseas.

Mr. K. Morgenstern scored the seedlings of this red spruce provenance experiment and supplemented this material with field observations. Using an Anderson hybrid index, he was able to show that the North Carolina red spruce is uniform and typical (black spruce is absent and its presence there during glaciation is doubtful). In Pennsylvania weak introgression occurs, resulting from the restriction of both species to small swamp communities. In the southern New England States, where there is often distinct ecological isolation, black spruce growing in the swamps and red spruce at higher elevation surrounded by hardwoods, there is only a slight infiltration of black into red spruce. In northern New York, Nova Scotia, New Brunswick and Quebec introgression is usually strong as the two spruces usually meet on upland sites.

NORWAY SPRUCE

The I.U.F.R.O. Norway spruce provenance experiment has been remeasured for comparison with other experiments of this series. The results will be of limited value only because of white pine weevil damage. The experiment was given a very heavy thinning removing weevilled and suppressed trees. Weevil attack has been studied in relation to measurements taken before weevil attack to see what influence form factors and provenance may have on weevilling. A number of plus trees have been selected and propagated to make family breedings.

An experiment comparing Finnish, Latvian, Polish, Russian and Siberian Norway spruce provenances with the fast growing German provenances from Swedish plantations and locally selected hardy material went on trail in 15 localities (Exp.No. 200). The Russian and Siberian Norway spruce are slower growing than most of the European Norway spruce grown here. However, the Russian and Siberian provenances may be of particular interest in the Boreal Forest.

· . · . . .

Open pollinated progenies of our apparently weevil resistant Norway spruce selections from Hydson's Place are on trial in New York and Ontario (Exp.No. 214). Our own experiment with this material (Exp.No. 120) was remeasured, but it is a little too early to assess weevil damage.

A number of <u>Picea abies x asperata</u> hybrids were planted in Ontario (Exp.No. 253). <u>Picea asperata</u> is not quite hardy at the Petawawa Forest Experiment Station and is badly weevilled. The hybrid may have some merit in areas with too warm summers for Norway spruce.

SPRUCE GRAFTING

An experiment with reciprocal grafting of white, red, black and Norway spruce gave interesting results. These four spruces were completely compatible, but scion growth was very much influenced by the rootstock. Black, Norway and white spruce scions grew poorly on red spruce rootstocks, while the black spruce rootstock stimulated scion growth of Norway and white spruce about 50 percent and red spruce scions by 130 percent. This stimulation (or lack thereof) may be explained in terms of efficiency of root systems and tolerance. Red spruce has an inefficient root system, has a high requirement for nutrition, and has the slow early growth typical of tolerant species. Elack spruce - on the other hand - seems to have an efficient root system, a low requirement for nutrition, and the rapid early growth typical of an intolerant pioneer species. White and Norway spruce are intermediate in these respects. These reciprocal grafts have been planted on four different sites to study their reaction to soil fertility and moisture regime. (Exp.No. 212)

RED PINE

A night frost in early June, 1960, severely damaged the red pine provenance experiments in the Young Creek Area (Exp. Nos. 38, 39 and 74-B). The north-western Ontario red pines showed less damage than local red pine.

A provenance experiment and a single tree progeny test of Ontario and Wisconsin red pine (Exp. Nos. 215 and 216) were followed with measurements.

Seed of single red pines from many different local stands were gathered partly for Michigan State University and partly to provide material to settle the problem of how best to sample a red pine stand (Exp. No. 238). If we handle only 50 seed lots, for example, should we take one tree from each of 50 stands, 5 trees from 10 stands, or 10 trees from 5 stands?

The morphology of red pine is remarkably uniform throughout its range. A few minor and very localized variations are known to exist, e.g., the fastigiate forms in Wisconsin and the "tassel pines" from Bancroft, Ontario. In general there is little or no basis for individual tree selection based on the phenotype (except perhaps in the isolated occurrences in northern Ontario where Dr. Heimburger has noticed genetic drift). Provenance hybridization is one possible way to increase variability in red pine.

In the spring of 1961 we co-operated with Mr. D.P. Fowler in performing controlled pollinations in three locations: (1) Petawawa Forest Experiment Station; (2) Watabeag Lake near Swastika; and (3) Lake Abitibi west of Cochrane.

At each locality, a selfing, a crossing to the same stand, and a State of Maine provenance crossing were made for Mr. Fowler on each of five trees. Our own project involved provenance hybridization using pollen from Pennsylvania, Massachusetts, New York, Maine, Michigan, Wisconsin and Minnesota. About one thousand bags were used.

Trouble was expected with the red pine cone borer (Conophthorus resinosae) and the trees were sprayed for two years with D.D.T. Even then the borer has taken a heavy toll in the Petawawa area, while there were no borers at Watabeag Lake.

JACK PINE

Jack pine is a tree breeder's delight. Its fast growth and early flowering makes for easy handling and rapid turnover.

In 1960 we went through our collection of jack pine grafts, pinched off all male flowers and left the females to be pollinated with whatever local jack pine pollen there was in the air. To insure adequate pollination we supplied extra pollen to the open flowers. With very little effort we were able to make the following provenance hybrids.

Somerset Co., Maine	
Cumberland Co., N.S.	
Northumberland Co., N.B.	
Lake St. John, P.Q.	
Opawica Distr., P.Q.	·
Laviolette-Letondal Co	P.Q.
Espanola, Ont.	
Crawford Co., Mich.	
Vilas Co., Wis.	
Sandilands. Man.	
Smith. Alta.	•
Fort Smith N.W.T.	2.1

x Petawawa, Ontario

A hybrid population from northern Alberta where the range of jack pine and lodgepole pine overlaps, was also crossed with Petawawa jack pine. The individuals were separated into the following four groups according to cone characteristics:

- 1) smooth closed cones

- 2) smooth open cones
 3) prickly closed cones
 4) prickly open cones

The individuals with most lodgepole characteristics have the best form, but not the best growth rate. Furthermore, a number of the lodgepole and jack pine hybrids from Placerville were backcrossed to Petawawa jack pine. These hybrids are of extremely poor form and are slower growing than local jack pine.

In 1961 jack pine provenance experiments and clone collections were used to make 43 different provenance hybrids. Northern provenances were crossed with southern, and eastern with western. About 900 bags were used.

One hundred jack pine seed lots have been assembled for an all range jack pine provenance experiment. The intention is to run a number of nursery provenance experiments in as many climates as possible using all - or nearly all provenances. We are trying a new technique with these experiments. Ordinarily we have planted 2-0 plants for our nursery provenance experiments. With the jack pine material we will sow seed spots on l' x l', use a 10-spot plot and up to 10 replications. The seed spots will eventually be weeded down to one plant per spot. The seed will be distributed to North Carolina, New Hampshire (three experiments), Massachusetts, Wisconsin, Michigan, New Brunswick, Quebec, and Longlac and Petawawa in Ontario. The same material will be grown in the growth chambers at Yale University. Other series of experiments using standard nursery practice and field planting will eventually be established across the jack pine range.

A number of older jack pine provenance experiments and observation plots have been measured twice to assess increment (Exp.Nos. 41, 41, 42 and 82). The Lake States jack pine provenance experiment that was planted in three locations in Canada has also been remeasured (Exp. Nos. 125, 132 and 133) and will be compared with the experiments planted in the Lake States.

Three hundred single tree progenies of jack pine will be studied for inheritance of cone opening, cone angle, branch angle and stem form. We found that jack pine from Massey, Ontario have a larger branch angle and more open cones (26%) than local jack pine. Petawawa jack pine had 16 percent open-coned trees and Notakim, Depot, P.Q., 11 percent.

SCOTS PINE

Our main effort has been the establishment of 24 provenance experiments and observation plots containing 13 Russian and Siberian Scots pine provenances (Exp.No. 201). Most of this material will be of particular interest in the Boreal Forest Region where hardiness counts more than growth rate. (Selected provenances from Europe should be faster growing than the Russian and Siberian provenances in the Great Lakes-St. Lawrence Forest Region). Two provenances from southern Russia (from Kozinsky, Kiev Forest District, Ukrania, and the Province of Woronesh) are of particular interest as potential sources for resistance to the European pine shoot moth, as they come from areas of high resin production.

Twenty-six new selections of potentially "perfect" Christmas trees were added to our Scots pine clone collection. Some of our earlier selections flowered and a limited number of controlled crosses were made. Lack of pollen on the young grafts was the limiting factor.

Observation plots of promising Scots pine provenances were established for future selection of Christmas trees. (Exp. No. 188). Seven Scottish, two French, one Hoch-Schwarzwald, one Campine, and one southern Ontario provenance went on trial. It is hoped that we will be able to make family breeding with this material. Such family breeding should give more uniform progenies than our earlier selections that have largely been of unknown parentage.

.

OTHER PINES

Single tree progenies of high and low resin yielders of <u>P. nigra</u> from Austria, several <u>P. nigra</u> varieties, and southern Russian Scots pine were planted in shoot moth tests on Turkey Point and at Elmira (Exp.No. 183). This is part of a co-operative study of European pine shoot moth preference, and of resistance breeding conducted by the Forest Research Branch and the Forest Entomology and Pathology Branch of the Department of Forestry, and the Ontario Department of Lands and Forests.

LARCH

Eleven provenance experiments and observation plots comparing <u>Larix</u> <u>laricina</u>, <u>L. kurilensis</u>, <u>L. decidua</u> and four Polish larch stands were planted in Canada and the New England States (Exp. No. 202).

Another experiment comparing European, Polish, Japanese, European x Japanese, and western larch was planted in 10 locations (Exp.No. 209). The experiments planted at Petawawa Forest Experiment Station were established along a moisture gradient to study the effect particularly on the European, the European x Japanese, and Japanese larch. It may well be that the Japanese larch and the hybrid will be superior on the moist sites while the European larch will grow best on the drier sites. Another interesting feature of this experiment was that two lots of <u>L</u>. <u>occidentalis</u> were fading away in the nursery and produced very small and poor plants. Our soil seems to lack some essential quality, possibly mycorrhiza.

OTHER SPECIES

Several birch experiments have been planted. Some are single tree experiments with paper birch and others are trials of exotic birches.

PLANT FORCING AND FLOWER INDUCTION

Quite a few research workers are working on techniques for accelerating growth. This is usually done by combining long days with adequate temperature and growing conditions. Several interesting differences between species have been found. However, sooner or later most plants stop shoot elongation by forming buds or become unhealthy because of too long exposure to an artificial environment. The spruces and the pines are particularly difficult to force. We have therefore concentrated on studying the chilling requirement of plants that have been treated with long days and long growing seasons. Preliminary experiments show that spruce chilled for four weeks will break dormancy only on the lower branches. Six weeks' chilling broke dormancy of the side buds, but not of the terminal bud. Only an eight weeks' chilling induced complete and normal flushing. We also tried chilling at different light intensities. Best was chilling in the dark (normal cold storage room). Low light intensity chilling in a growth chamber gave intermediate results. Poorest was chilling at high light intensity. Plans for experiments with multiple growth cycles in one year, giving cycles of long days and warmth, short cool days, and a short period of chilling have been planned, but not done as Mr. Yeatman left for Yale.

A number of branch girdling experiments were done on spruce and pine in an attempt to pin down when the flower premordia are formed. Many of the branches broke and some of the experiments were not conclusive.

In red pine the girdling at any date reduced female flowering to about one third. Male flowers were stimulated 61 to 76 percent on branches girdled up till June 1; with June 15th girdling, there was a 7 percent reduction and on August 1 a 35 percent reduction.

In jack pine, female flowering is stimulated by girdling until July 1, while the male flower stimulation stopped with girdling on May 15.

A timing experiment with pruning and ammonium fertilizer on young red pine gave indications of how one may treat red pine seed orchards. Any kind of pruning inhibits female flowering (as might be expected). Pinching of lower side branches in early May promotes male flowering, but this effect disappears when the trees are fertilized. To reduce growth and increase branching the leader and upper whorls should be pruned between June 9 and July 9. This will triple the number of branches and, as the branches are smaller, should eventually be conducive to female flowering, and later also to male flowering (one only finds male flowers on small branches). The lack of male flowers in seed orchards, and in the tree breeders clone collections could be overcome by pinching of the lower branches in early May.

In white, black and Norway spruce, branch girdling was a complete failure as the incidence of breakage was very high. The timing experiment was therefore repeated using treatment of whole trees (girdling, strangulation and rootpruning). For 16-year-old white spruce and 15-year-old Norway spruce only rootpruning before May 10 was effective. Black spruce reacts to treatment up to June 15 (although early treatments are best) and reacts better to girdling and strangulation than white and Norway spruce.

Six-year-old white spruces were treated with a combination of short day, high lime, and spraying the uracil, xanthine and ethylenechlorhydride did not produce any flowers. Man-high red pines sprayed with uracil and xanthine did not flower either, but 2-year-old larch seedlings sprayed with uracil yielded 10 percent female flowers.

Mr. M.M. Giertych made a comprehensive study of red pine buds, comparing different age classes, different positions within the crown, and ammonium nitrate-fertilized and non-fertilized trees. Three growth promoters and one growth inhibitor were found. Terminal buds had higher levels of growth promoters than side buds indicating a hormone gradient in flowering. The effect of the fertilizer was to increase the content of all growth regulators, particularly the inhibitor which may explain the reduction of leader growth seen on red pine fertilized with ammonium nitrate. DNA concentration was proportional to number of cells and therefore constant per cell, while RNA varied with season and bud position. The ammonium nitrate fertilizer increased the RNA concentration in July. This is a very important study for our understanding of the physiology of flowering - but we still have a long way to go and we need more of these fundamental studies.

VEGETATIVE PROPAGATION

Some of the spruce grafting experiments have been mentioned above.

Red pines grafted on Scots pine rootstocks show incompatibility. Following are typical percentages for red pine grafted out of doors on large Scots pine rootstocks. The first year survival of about 1,000 grafts was 72 percent. Eight years after we had the following:

Percent of first year survivors

54

: 5

Healthy green Yellow needles Burned needles Dead recently with brown needles Dead (reason suspected to be incompatibility)

Total:

100

Thirty-six percent have died. Another 10 percent are unthrifty with yellow or brown needles typical of starved foliage.

A total of 3,788 grafts were made during the two-year period. Of particular interest are the <u>Pinus nigra x resinosa</u> hybrids from P**lacerville** (Table 1).

NURSERY WORK

The nursery has been enlarged, but it will be some time before the new area is uniform and of adequate fertility. Many new Dunnemann beds have been installed. About 125,000 seedlings were transplanted (Table 2).

a de la companya de l La companya de la comp

PLANTATION WORK

Thirty-nine experiments including 73,383 plants covering 39.4 acres were field planted at the Petawawa Forest Experiment Station. About 2,500 of these were grafts (Table 3).

Another 72 experiments were planted by co-operators in Canada and the United States. These included 181,477 plants and covered 163.2 acres (Table 4).

About 29,000 plants were distributed for another 7 experiments to be field planted in 1962 (Table 5).

An increasing amount of time is spent looking after older plantings and in getting new experiments properly established.

Table 1

SUMMARY OF GRAFTING DURING SPRING AND FALI	1960 AND 1	<u>961</u>	
1. SPRING 1960	•		
<u>P-139 & P-142, Exp. No. 128</u>	No. Grafts		* .
17 clones of <u>P.resinosa</u> from 3 areas in northern Ontario.	660		
<u>P-133</u>	- 1		
5 clones of <u>P.glauca</u> from Petawawa	240		
<u>P-60. Exp. No. 241a; P-133</u> 4 clones of P.glauca from Petawawa		·	
(grafting and pruning experiment)	560	The second	
Exp. No. 248 Cold storage of red pine, white spruce &			
Norway spruce	300		
		1,760	
2. FALL 1960			
P-139 & P-142. Exp. No. 238		,	
43 clones of <u>P. resinosa</u> from Petawawa F.E.S.	422		
P-139 & P-142, Exp. No. 257	· .		
15 clones of <u>P.resinosa</u> from 14 provenances in Ontario	130	n an	÷ .
6 clones of <u>P</u> . sylvestris x <u>P.mugo</u> F_3	60	612	
Total grafting 1960	. × ¹	2,372	х
3. SPRING 1961			
<u>P-144</u>	n an an Al Al Anna an Al	:	
2 clones of <u>P. nigra x resinosa</u> & the parent trees	14		*
<u>P-147</u>			
10 clones and 2 mixtures of clones of <u>P.sylvestris</u> (Christmas tree selections)	255 w 1.1		
P-133			
4 clones of <u>P.glauca</u> from Causapscal, P.Q. and Petawawa, Ontario	195		

.M-11

5 • • • •

Table 1 (Cont'd)

4 clones of	<u>241b; P-133</u> <u>P.glauca</u> from Petawawa		
(Signering a	na pruning experiment)	480	944
-137			
l clone and species from 6 clones of	l population of exotic spruce m China P.jezoensis hondoensis x glau	<u>ca</u> 33	· · ·
l clone of surrounding	P.glauca x mariana & 8 clones of white and black spruces	of 69	
	an a		1,076
147	4. FALL 1960		n jan
14 clones of (Christmas f	f <u>P.sylvestris</u> tree selection)	340	•
1. N. 18		• .) • •	340
÷	Total grafting 1961		1,416
e estador e construir e estador de la construir			
	$\{1, \dots, k\} = \sum_{i=1}^{n} (1 - i) \sum_{i=1}^{n} $		·: · ·
		n an	
•			a.
· · · ·		a an	• • • •
· · · ·	· · · ·		
	الم الألم المعالية (من المعالية المعالية المعالية المعالية المعالية المعالية المعالية المعالية المعالية المعال الم أولية في المعالية (من المعالية المعالية المعالية المعالية المعالية المعالية المعالية المعالية المعالية المع		

and the second second

-

Table 2

TRANSPLANTING OF TREE BREEDING MATERIAL

1. SPRING 1960

Project or		
Exp. No.	Material	No. Planted
216	<u>Pinus resinosa</u> provenances from Lake States and eastern Canada	26,000
215	<u>Pinus resinosa</u> , single tree progenies from Lake States and Grand Bend, Ont.	12,500
P-147	<u>Pinus sylvestris</u> crossing for Christmas tree breeding	60
P-147	<u>Pinus sylvestris</u> of different origin for Christmas tree selection	3,170
183-E	<u>P. nigra austriaca</u> provenances from Yugoslavia	3,000
218	<u>Picea glauca</u> provenances from northern Canada (from Labrador to British Columbia) <u>Betula verrucosa</u> , elite tree from Sweden Pine and spruce rootstocks	1,250 160 6,000
		- and the second second second
	Total transplanting 1960	52,140
	2. SPRING 1961	
194	<u>Picea glauca</u> provenance experiment	66,000
263	<u>Pinus sylvestris</u> provenances	2,600
264	<u>Pinus nigra</u> provenances	700
35B_C	Larix species and species hybrids	1,400
	Picea " " "	1,520
	<u>Betula papyrifera</u> single tree and popu- lation collections from Petawawa	1,000

Total transplanting 1960

73,220

Table 3

وهيد وران والمراجع المراجع

SPRING PLANTING 1960 AND 1961 BY THE TREE BREEDING SECTION PETAWAWA FOREST EXPERIMENT STATION

and an and an area

the second second second second

Exp.		No. of	Area	Plantation
.10.	Description	Plants	Acres	Area
	Seed Plants 1960			en e
95-H	Provenance experiment including 14 races of red spruce.	7760	2.85	P.A.114
01-0	Trial of Scandinavian elite birch for future breeding purposes.	637	0.35	P.A.106
39	Comparison of Alaska sitka spruce, white spruce and sitka x white	3482	1.28	P.A.114
	spruce hybrids, Danish white spruce and sitka x white spruce hybrids from plantations in Jutland and fro controlled hybrids in Horsholm; Cochrane white x B.C. sitka and			
	local white spruce.		· ·	
38 _ A	Selection of Christmas trees in Schwarzwald, Scottish and south- ern Ontario Scots pine provenances.	9852	3.62	P.A.120
)1-A	Provenance experiment and obser- vation plots including Russian and Siberian Scots pine.	6663	2.45	P.A.121
)1 _ B	Ditto	5400	1.98	P.A. 120
)2 _ A	Provenance experiment and obser- vation plots including various larch species.	3999	3131	P.A.122
)2 _ J	Ditto	5159	4.26	P.A.114
	Total 1960	42,952	20.10	· · · ·
	Seed Plants 1961			
open :	One-parent progeny test with red spruce, red x black spruce hy-	79	0.11	P.A.103

M-14	4
------	---

1.1

Table 3 (Cont'd)

Exp. No.	Description	No. of Plants	Area Acres	Plantation Area
119 <u>open</u>	Transfer of weevil resistance from white spruce to Norway spruce	490	0.40	P.A.104
	by means of crossing bridges, white sitka-Norway spruce, and others.	- 1		
119 <u>open</u>	Ditto	450	0.76	P.A.116
200 – J	Provenance experiment and obser- vation plots including north Euro- pean and Russian Norway spruce.	3082	1.13	P.A.123
200 - K	Ditto	6850	2.51	P.A.114
202 - J	Provenance experiment and obser- vation plots including various larch species.	5159	4,26	P.A.114
208 <u>open</u>	Comparison of good and poor geno- types in white spruce.	12	0.04	P.A.104
209 - A	Provenance experiments and obser- vation plots including <u>Larix decidua</u> <u>L. leptolepis</u> , <u>L. eurolepis</u> and <u>L.</u> <u>occidentalis</u> .	, 3793 ,	1.40	P.A.123
209 - F	Ditto	1150	0.95	P.A.123
209-F-1	Ditto	1657	0.69	P.A.124
209-F-2	Ditto	836	0.35	P.A.124
209-Н	Ditto	164	0.24	P.A.114
217 - A	Co-operative study of Canadian, European, Korean and Japanese birches.	990	0.44	P.A.123
217-B	Ditto	105	0.09	P.A.101
217 - C	Ditto	1800	1.49	P.A.124
217-D	Ditto	403	0.33	P.A.106
243 <u>open</u>	Interspecific hybridization in Picea	13	0.04	P.A.104

. .

Table 3 (Cont'd)

.

Exp. No.	Description	No. of Plants	Area Acres	Plantation Area
253 - A	<u>Picea abies x asperata</u> hybrids suitable for warm summers.	119	0.20	P.A.116
2 53 - B	Ditto	780	0.64	P.A.114
254 <u>open</u>	Interspecific hybridization in <u>Abies</u> .	23	0.03	P.A.104
	Total 1961	27,955	16.10	
	Total 1960	42,952	20.10	
	Seed Plants, Grand Total 1960-61	70,907	36.20	
	Grafts 1960			
87 <u>open</u>	Investigation of jack pine, lodgepole pine and their hy- brids for resistance to sweet fern blister rust.	1278	0.23	.P.A.106
128 <u>open</u>	Grafted red pine population sample intended for provenance hybridi- zation.	s 148	0.12	P.A.104
	Total 1960	 1426	0.35	
	<u>Grafts 1961</u>	1720	0.))	
36 <u>open</u>	Selection of the perfect Scots pine Christmas tree.	105	0.34	P.A.115
37 <u>open</u>	Investigation of the jack pine, lodgepole pine and their hybrids for resistance to sweet fern blister rust.	5	0.01	P.A.115
12-A	Interspecific grafting of black, Norway, red and white spruce.	320	1.06	P.A.115
12 - B	Ditto	160	0.53	P.A.116
12_C	Ditto	80	0.07	Nurserv

, ;

. . .

Table	3	(Cont	'd)

Exp. No.	Description	No. of Plants	Area Acres	Plantation Area
212-D	Interspecific grafting of black, Norway, red and white spruce.	80	0.07	Nursery II
242 <u>open</u>	Selection of white spruce plus trees.	126	0.19	P.A.114
:42 <u>open</u>	Selection of white spruce plus trees.	160	0.53	P.A.116
56 <u>open</u>	Grafted white spruce population samples.	14	0.05	P.A.116
. ⁴	Total 1961	1,050	2.85	
· .	Total 1960	1,426	0.35	
	Grafts Grand Total 1960-61	2,476	3.20	
	Seed Plants Total 1960-61	70,907	36.20	
	ta an	73,383	39.40	• :: • • • : ::

 $(\mathbf{q}_{1}) \in [\mathbf{q}_{2}]$

i

1.1۱Ì

•1 ; Ę. ŧ

· · · .

• .:

 $\tilde{\mathcal{L}} \sim \sqrt{L}$ a starter and Bertheline groups with the barrenge track 19. <u>19.</u> 19.

Table 4

		•	• • •	
Sxp. No.	Title	No. of Plants	Area Aores	Agency & Location
	Seed Plants 1960			
1 C	Comparison of wood production and growth behaviour of 18 se- lect broad and slender white spruces of various provenances by means of single tree progeny	7 <i>9</i> 8	0.29	J.Blimkie, Chalk River, Ontario
	test.			
2-0	Single tree progeny test in- cluding 16 P.F.E.S. white spruces	1102	0.40	J.Blimkie, Chalk River, Ontario
3-G-3-1	Ditto	144	0.12	Mich. State Univ., East Lansing, Mich., U.S.A.
3-6-3-2	Ditto	660	0•54	Mich.State Univ., East Lansing, Mich., U.S.A.
3-M-1	Ditto	708	0.26	J.Blimkie, Chalk River, Ontario
3-M-2	Ditto	405	0.15	J. Blimie, Chalk River, Ontario
5-B	Provenance experiment includ- ing 22 races of red spruce.	10,349	3.80	Dept. of Forestry, Acadia F.E.S., N.B.
5 - C	Ditto	10,481	3.85	Dept. of Forestry, Fundy National Park, N.B.
5-K	Ditto	700	1.00	Lake States F.E.S.,
		n an thair Chuir an thair Chuir an thair		Aninelander, Wisc., U.S.A.
-L-1	Ditto	900	0.51	New Hampshire For- est & Recreation
*		مربع بر ا	n an statu a statu ga statu	Committee, Concord, N.H., U.S.A.

Table 4	(Cont'd)

Exp.			No of	Area	and the second
No.	Title		Plant	s Acres	Agency & Location
95-L-2	Frovenance experiment including 22 races of red spruce.	1–.	1,100	0.63	New Hampshire Forest & Recreation Commis- sion, Concord, N.H., U.S.A.
95-n	Ditto		2,019	1.00	Univ. of Toronto, Ranger School, Dorset, Ont.
95-0	Ditto		1,750	2.00	Mr. S. Davis, Eganville, Ontario
95-P	Ditto	· ·	1,400	1.15	Ont. Dept. of L. & F., Sault Ste. Marie, Ontario.
95-Q	Ditto		1,669	1,40	Univ. of New Bruns- wick, Maritime For. Ranger School, New Maryland, N.B.
95-R	Ditto		7,000	5.80	Acadia Univ., Wolf- ville, N.S.
95-S	Ditto		9,425	8,40	Bowater's Mersey Paper Co., Liver- pool, N.S.
95-T	Ditto	•	336	0.12	Dept. of Forestry, For.Res.Br., Fredericton, N.B.
95-U	Ditto		4,700	5.00	Nova Scotia Dept.of L. & F., Yarmouth, N.S.
L83-B-2	Testing of single tree pro- genies of high resin yield- ing Austrian pine for shoot moth resistance.		150	0.12	Dept. of Forestry, Forest Insect Lab., Sault Ste. Marie, Ontario.
.88 - B	Selection of Christmas trees in Schwarzwald, Scottish and southern Ontario Scots pine provenances.	:	3,820	0.62	Mr. J. Blimkie, Chalk River, Ont.

M-	1	9

Table 4 (Cont'd)

·· · .

Exp.	Title	No. of Planta	Area	Aconor & Location	
1.0.	*1 616	I Lan US	NOTES	ngency a nocation	· · ·
200-D	Provenance experiment & ob- servation plots including north European & Russian Norway spruce.	3,298+ local jp	4.00	Spruce Falls Power & Paper Co., Kapus- kasing, Ontario	
200-G	Ditto	910	0.75	Univ. of Toronto, Faculty of Forestry, Toronto. Ontario.	•• <u>·</u>
. 1				-0101109, 01100-201	
201-C	Provenance experiment and ob- servation plots including Russian & Siberian Scots pine.	4,096	3.40	The KVP Co., Espanola, Ontario.	··· .·
2 01-D	Ditto	4,096	3.40	Kimberly-Clark Pulp & Paper Co., Longlac, Ontario.	
201-E	Ditto	3,650	3•75	Univ. of Toronto, Ranger School, Dorset, Ontario.	·
201-F-1	La Ditto	3,185	2.63	Dept. of Forestry, Manitoba.	
201-F-2	Ditto	3,185	2.63	Dept. of Forestry, Manitoba.	
?01 - K	Ditto	2,020	2.30	Forest Nursery Sta- tion, Indian Head, Sask.	
201 - H	Ditto	4,088	1.51	Sask.Dept. of Nat. Res., Forestry Br., Prince Albert, Sask.	*,**
201-1-1	Ditto	792	3.50	U.S. Forest Service, North.Inst. of For. Genetics, Rhine- lander, Wisc., USA.	
201-I-2	Ditto	792	3.50	U.S. Forest Service, North.Inst. of For. Genetics, Rhine- lander, Wisc., USA.	
M-20

Table 4 (Cont'd)

Exp.	M (), n	ne odrahom za naj kole na naj za si	No. of	Area	
MO.	Title		Plants	Acres	Agency & Location
201-I-3	Provenance experiment and servation plots including Russian & Siberian Scots	ob- pine.	792	3.50	Forest Nursery Sta., Indian Head, Sask.
201-I-4	Ditto		792	3.50	U.S. Forest Service, North.Inst. of For. Genetics, Rhine- lander, Wisc., USA.
201-I-5	Ditto		792	3.50	U.S. Forest Service,
					Genetics, Rhine- lander, Wisc., USA.
201 -J	Ditto		770	0.20	Univ. of Toronto, Faculty of Forestry
201-K	Ditto		3,102	2.50	Ont.Dept. of Lands & Forests, Maple, Ontario.
201-L	Ditto		200	0.16	Forest Insect Lab., Sault Ste. Marie, Ontario.
202-0	Provenance experiment and servationpplots including various larch species.	ob-	3,400	3.00	Bowater's Mersey Paper Co., Darling Fields, N.S.
2 02-D	Ditto		670	0.55	Univ. of Toronto, Faculty of For., Toronto, Ontario.
202 - F	Ditto		120	0.10	Univ. of New Bruns- wick, Fredericton, N.B.
202-G-1	Ditto	2	2,400	3.50	Can.Int.Paper Co., Harrington For. Farm, P.Q.
2 02-G-2	Ditto	. ε	3,102 11	L•84	Can.Int.Paper Co., Harrington For. Farm, P.Q.

M-21

Table:	4	(Cor	t'd)	e . j
				·

۰.

• •

Exp.	an a		No. of	Area	(4) A set of the se
No.	Title	ء محمد محمد محمد محمد محمد محمد محمد محمد	Plants	Acres	Agency & Location
202-I	Proversace experim servation plots i various larch spe	ent and ob- ncluding cies.	650	0. 50	S. Davis, Eganville, Ontario.
2 02 - K	Ditto	al an i	1,900	1.57	James McLaren Co., Buckingham, P.Q.
209-B-1	Provenance experimentary servation plots in <u>decidua</u> , <u>L. lepto</u> <u>occidentalis</u> and <u>lepto</u>	ments and ob- ncl. <u>Larix</u> <u>lepsis, L</u> . <u>L. eurolepsis</u> .	2,700	4.00	Can.Int.Paper Co., Harrington For. Farm, P.Q.
209-B-2	Ditto		6, 500	9.50	Can. Int. Paper Co., Harrington For. Farm, P.Q.
209 - C	Ditto		770	0.28	Univ. of Toronto, Faculty of Forestry, Toronto, Ontario.
n yes en e Nise Nise	Seed Plan	ts 1960 Total Seed Plants 196	123,388	116.7	3
183-,D	Testing of single high resin yield A shoot moth resists	tree progenies ustrian pine f unce.	of 360 or	0.20	Ontario Dept. of L. & F., Maple, Ont.
200-A	Provenance experim vation plots inclu European and Russi	ent and obser- ding north an Norway sprue	60 ce.	0.05	Univ. of New Bruns- wick, Fredericton, N.B.
200-F	Ditto		3,781	3.12	Sask.Dept, of Nat. Res., Forestry Br., Prince Albert, Sask.
200-L	Ditto	en e	2,000	1.15	La Maison Provincial des Freres Maristes, Levis, P.Q.
200–M	Ditto		2,000	1.15	Leopold Dion, Belle- chase, P.Q.

•

с. •

5411

المعالم والعار

1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 -		
at a terr	M-22	
	Table 4 (Cont'd)	

Exp.		No. of	Area	· · · · · · · · · · · · · · · · · · ·
No.	Title	Plants	Acre	s Agency & Location
201 - M	Provenance experiment and ob- servation plots including north European and Russian Norway spruce.	3,600+ 400 of loca materia	3.30 1 1	Ontario Dept. of L. & F., Swastika For- est District, Ont.
201 - N	Ditto	3,600+ 400 of loca materia	3.30 1 1	Ontario Dept. of L. & F., Swastika For- est District, Ont.
201-0	Ditto	197	0.37	Dept. N.A.& N.R., North. Adm. Branch, For. Exp. Nursery, Fort Smith,,N.W.T.
201 - P	Ditto	72	0.11	La Maison Provin- cial des Freres Maristes, Levis, PQ.
201-Q	Ditto	560	0.50	Mr. S. Davis, Eganville, Ontario.
2 01-R	Ditto	2,565	0.61	Mr. J. Blimkie, Chalk River, Ont.
201 - S	Ditto	60	0.05	Univ. of New Bruns- wick, Fredericton, N.B.
202-E	Provenance experiment and ob- servation plots including various larch species.	132	0.30	Dept.of Forestry, Acadia F.E.S., N.B.
2 09 - D	Provenance experiments and ob- servation plots incl. Larix decidua, L. leptolepsis, L. eurolepsis, & L. occidentalia	1,150	1.68	Dept. of Forestry, Acadia F.E.S., N.B.
ы. н а	<u>earorepois</u> , w <u>s</u> . <u>occuentaris</u> .	901	1.33	
209 - G	Ditto	140	0.21	La Maison Provincial des Freres Maristes, Levis, P.Q.
214-A	Trial of Norway spruce prove- nances & frost hardy and weevil rated Norway spruce selections from Hudson's Place Planted in <u>Pack</u> Forest	441	0.50	N.Y. State Univ. Coll. For., Syracuse, N.Y., USA.

M-23	
------	--

•

	r=2 5	
an a	Table 4 (Cont'd)	

6,1964	· · · · ·		<u></u>	<u></u>	
Exp.	"		No. of	f Area	
NO.		+	Plants	Acres	Agency & Location
214 - B	Trial of Norway spranances & frost hardy rated Norway spruce from Hudson's Place Planted in <u>Pack</u> Fore	ace prove- 7 and weevil selections est	441	0,50	N.Y. State Univ. Coll For., Syracuse, N.Y., U.S.A.
214-0	Ditto		441	0.50	N.Y. State Univ. Coll For., Syracuse, N.Y., U.S.A.
214 - D	Ditto		25,300	. 21.0	Ont. Paper Co., Gore Bay, Ontario.
253-0	Suitability of <u>Pices</u> <u>asperata</u> hybrids for summers.	<u>abies</u> x warm	1,200	1.00	Ont. Paper Co., Wanifleet Twp., Ont.
253-D	Ditto	$M_{\rm eff} = M_{\rm eff} + \frac{1}{2} M_{\rm eff}$	7,000	4.0	R. Wagner, Courtland, Ontario.
253 - E	Ditto		1,000	0.68	La Maison Provincial des Freres Maristes, Levis, P.Q.
253 - F	Ditto	· · · · ·	1,000	0.57	Leopold Dion, Belle- chase, P.Q.
• * * *	$c = C^{+}$				
	ta ang ta ang Ta ang ta ang	otal 1961	58,001	. 46.18	3
	T	otal 1960	123,388	116.73	
	Seed Plants	1960-61	181,389	162.91	
		Grafts 196	1		
86- <u>open</u>	Selection of the per pine Christmas tree.	fect Scots	88	0.29	W.E. Nodwell, Court- land, Ontario.
				0.29	
Total :	number of plants			2	
plante 1960-63	l by cutside agencies l Gr Se	afts ed Plants	88 181,389	.29 162.91	
			181,477	163.20	

M-24

DISTRIBUTION OF PLANTS TO OUTSIDE AGENCIES IN 1960 AND 1961

(Probably field planted in 1962)

Exp. No.	Agent and Planting Site	No. of Seedlots	No. of Seedlings
200 - B	Dept. of Forestry, Manitoba, Riding Mountain Nursery. (Norway spruce, 2-1 stock)	12	7200
200 -C	Dept. of Agriculture Forest Nursery Station, Indian Head, Sask. (Norway spruce, 2-1 stock)	11	6900
200-E	Kimberly-Clark Pulp & Paper Co., Longlac, Ont. (Norway spruce, 2-1 stock)	6	7000
200-Н	Northern Institute of Forest Genetics, Rhinelander, Wisc., U.S.A., (Norway spruce, 3-0 stock)	6	3000
1-00 - 1	Univ. of Minnesota, St. Paul, Minn. (Norway spruce, 3-0 stock)	5	2500
00 N	Dept. of Northern Affairs & National Resources, Northern Administration Branch, Fort Smith, NWT. (Norway spruce, 2-2 stock)	12	300
01-G	Dept. of Agriculture, Forest Nursery Station Indian Head, Sask., Scots Pine, 2-1 stock.	, 11	2200
	Total:	63	29,100

PUBLICATIONS

- HOLST, M. J. Seed selection and tree breeding in Canada. Paper prepared for the 8th British Commonwealth Forestry Conference, East Africa, 1962. For. Res. Br. Tech. Note No. 115. pp. 34
- MORGENSTERN, E.K. and YEATMAN, C.W., Editors. Proceedings of the 7th Meeting of the Committee on Forest Tree Breeding in Canada, Lake Cowichan, B.C., Aug. 21-23, 1960. Forest Research Branch, Canada Department of Forestry, Ottawa.

REPORTS

- HOLST, M. J. Co-operative white spruce provenance experiment for the Great Lakes-St. Lawrence Forest Region. Progress Report prepared for Forestry Committee, Woodlands Section, Canadian Pulp and Paper Association, Montreal, November 21, 1961. pp. 8.
- HOLST, M. J. Breeding of men and trees. Lecture. Deep River Science Association.
- HOLST, M. J. and SANTON, J.B. Reciprocal scion-rootstock combinations of <u>Pinus banksiana</u> Lamb., <u>P. resinosa</u> Ait., and <u>P. sylves-</u> <u>tris</u> L. pp. 30.
- MORGENSTERN, E.K. Planting in 1961 by the tree breeding section, Petawawa Forest Experiment Station. pp. 9.
- MORGENSTERN, E.K. The origin, early development and possible genetic background of the Hudson's Place Norway spruce. Exp. No. 4. pp. 9
- YEATMAN, C.W. Controlled pollination of black spruce in Manitoba, 1961. P-135, Exp. No. 243. pp. 10.
- YEATMAN, C.W. Controlled pollination of red pine, 1961. (Petawawa Forest Experiment Station and Watabeag Lake, Swastika) P-139, Exp. No. 128. pp. 15

M-25

REPORT ON TREE IMPROVEMENT AT THE ACADIA FOREST EXPERIMENT STATION, April, 1960 - March, 1962

by

H. G. MacGillivray

This report deals with tree improvement by the Maritimes District, Department of Forestry, from June 1960 to March 1962. This was concerned mostly with provenance studies of <u>Abies balsamea</u> (L.) Mill., <u>Picea abies</u> (L.) Karst., <u>P. rubens</u> Sarg., and <u>Pinus resinosa</u> Ait. Work was also done on <u>P. sitchensis</u> (Bong.) Carr., and several exotic <u>Larix</u> Mill. and <u>Abies</u> Mill.

A provenance test involving 17 seed sources of <u>Abies balsamea</u> was established (Fig. 1 and Table 1). Plantations A, B and C were planted by the Department of Forestry, and the others by co-operators. Plantation N, a direct sowing made in 1957 on cultivated land, was a failure.

In co-operation with the Newfoundland District, a study is being made to determine whether heredity is a factor in the (reported) premature loss of vigor of <u>A</u>. <u>balsamea</u> growing on the Burin, Avalon, and Bonavista Peninsulas of Newfoundland, here referred to as the "problem area". The planting stock, from seed collected in this and other areas in Newfoundland, and in New Brunswick, was grown in the Acadia Nursery. It was sprayed with insecticide in the autumn of 1960 and spring of 1961 to prevent the carrying of <u>Adelges piceae</u> (Ratz.) into uninfested areas of Newfoundland.

¹Research Officer, Forest Research Branch, Department of Forestry, Fredericton, New Brunswick.



Fig. 1. Locations of plantations in Abies balsamea provenance test

Plantation

Place

Salmon River Burn, Victoria County, New Brunswick. A В Acadia Forest Experiment Station, New Brunswick McDonald Brook, Restigouche County, New Brunswick. C D Haut St. Isidore, Gloucester County, New Brunswick. E Riley Brook, Victoria County, New Brunswick. F South Nelson, Northumberland County, New Brunswick. G Kingsclear, York County, New Brunswick. Lake George, York County, New Brunswick. H Ι Nasonworth, York County, New Brunswick. Riverside, Albert County, New Brunswick. Waweig, Charlotte County, New Brunswick. J K L Tynemouth Creek, St. John County, New Brunswick. Near Oxford, Nova Scotia. M St. Martine, St. John County, New Brunswick. N

Table 1. ABLES BALSAMEA PROVENANCE TEST

. .

Critician and Cr	Numbe	er of '	frees	Planted S	Sprin	g 1961.	Direct Sowing Spring 1957
Origin of Seed	<u>A</u>	<u>_B</u> _	<u>_C</u>	<u>D to</u>	<u>1²/</u>	M	<u>N</u>
Green River, N. B.	300	100	121	300	(J)	100	. . .
Oromocto, N. B.	1588	3/ ₁₀₀	121	700	(J)	100	1/2 lb.
Acadia Forest Experi- ment Station, N. B. (Seed lot MS. 118-56)	300	100	121	1400	(E)	100	1/2 lb.
Marystown, Burin Peninsula, Nfld.	-	100	8 12	-	-	100	
Sandy Brook, Nfld.	300	100	121	1000	(F)	140	1 1b.
Valcartier, P. Q.	300	100	121	200	(I)	100	<u> </u>
Salmonier, Avalon Peninsula, Nfld.	300	100	-			100	
Hawke's Bay, Nfld.	300	100	121	800	(I)	350	.
Bonne Bay, Nfld.	300	100	121	400	(L)	100	-
Nutak, Labrador	-	3			8		
Canada Bay, Nfld.	-	100	121	500	(K)	200	—
Duck Mt., Man.	300	100	121	æ.	-	350	-
Clear Lake, Man.	300	100	121	900	(L)	350	
West Canada Bay, Nfld.	-	100	420			50	-
White Bay, Nfld.	300	100	121	-		450	-
Adirondack Mts., N. Y.	300	100	121	600	(K)	100	-
Acadia Forest Experiment Station, N. B. (Seed lct MS. 304-56)	-	100	121	1000 2000 1000	(D) (G) (H)	200	-
TOTAL	4888	1603	1573	10800		2900	
spacing:	6' x 6'	4" x 4"	6.6x 6.6'	5' x 5' or 6' x 6'			

1/ Design: 3 x 4 rectangular lattice with three replications, and 100 trees per plot.

2/ Letters in brackets indicate plantation.

3/ Three hundred plot trees plus 1288 division and surround trees.

Plantation A on the Avalon Peninsula, within the "problem area and Plantation B in an area producing good <u>A. balsamea</u>, were planted by personnel of the Newfoundland District under a separate project. Plantations C, D and E were established by the staff of the Maritimes District, and F and G by co-operators (Fig. 2 and Table 2).

<u>A. balsamea, A. ernestii</u> Rehd., <u>A. homolepis</u> Sieb. and Zucc., <u>A. nordanniana</u> (Steven) Spach, <u>A. koreana</u> Wils., <u>A. koreana x veitchii</u> Lindl., and X <u>A. vilmorinii</u> Mast. planted at the Acadia Forest Experiment Station during the second week of June 1959 (1930 trees in all), and <u>A.</u> <u>balsamea, A. koreana, A. grandis</u> (Dougl.) Lindl., <u>A. concolor</u> (Gord. and Glend.) Lindl. and <u>A. alba</u> Mill. planted in the second week of June 1960 (4,150 trees in all), were inspected in 1961. Survival of <u>A. balsamea</u> was only 66 per cent in the 1959 planting and 51 per cent in the 1960 planting. The survival of the exotics was better than the <u>A. balsamea</u> except for the <u>A. homolepis</u> and <u>A. koreana</u> in the 1959 planting and the <u>A. grandis</u> in the 1960 planting. The low survival may have been caused in part by the lateness in the seasons when they were planted.

A provenance test involving 12 seed sources of <u>Picea abies</u> was established in the spring of 1961. An additional plantation was made in 1962 (Fig. 3 and Table 3). Eleven of the seed lots were provided by the Petawawa Seed Bank and originated in some of the best stands in Europe. Plantations A, B, Cl and C2 were planted by the Department of Forestry and the rest by co-operators. <u>P. glauca</u> (Moench) Voss was used in control plots in most plantations. <u>P. abies</u> planted by others at earlier dates in a number of areas in the Maritimes, appears to grow more rapidly than our native <u>Picea</u>. The plantations in this provenance test are located in a wide range of climatic zones and sites. Plantations A, B, G, H, J, K, L and most of the plots in I were visited in the Autumn, 1961. Little mortality was observed.

Plantation I consists of a number of plots in the foggy climatic region in the vicinity of Advocate, Nova Scotia. It was intended that these plots would be located on the slopes of the Cobequid Mountains, but some were planted on the outwash soils near the shore. The plots on the slopes appear more vigorous than the latter. Plantation J suffers from severe competition from hay.

The purposes for planting <u>Picea sitchensis</u>, <u>P. glauca</u> and lots likely to contain hybrids of these species, are to grow material, especially <u>P. sitchensis x glauca</u> hybrids, for use in the <u>Picea</u> A. Dietr. breeding at Petawawa, and to see how these lots grow in east-coast areas (Fig.4 and Table 4). Plantations B, C and D were established by co-operators. The direct sowing in 1957, Plantation D, was a failure.

The condition of each tree in Plantations A, B and C of the <u>Picea rubens</u> provenance test planted in 1959 and 1960 was recorded (Table 5). The foliage of trees in Plantations B and C on old fields was not as green as that in Plantation A, on a clear-cut area. Foliage on unreplicated observation plots in New Brunswick and Nova Scotia also showed this relationship.

Examination in June, 1960, of the <u>Pinus resinosa</u> provenance test planted in 1958 (4 x 4 balanced lattice square design with 16 seed sources, five replications, 49 trees per plot, plus division and surround trees, 5609 in all) showed heavy damage by deer; 22 per cent were browsed lightly to moderately, 73 per cent heavily and two per cent so severely that no buds remained to produce new shoots in 1961. Three per cent were dead. In the autumn of 1960 the terminal buds on the leaders which would produce the straightest stems were covered with polyethylene film to protect them from browsing deer, using a method described by McNeel and Kennedy (1959). The polyethylene was removed in the spring.



Figure 2. Locations of plantations in the study of premature loss of vigour of <u>Abies balsamea</u>

Plantation	Place
A	Near Salmonier Avalon Peninsula, Newfoundland.
В	Near Cormack, Newfoundland.
C & D	Salmon River Burn, Victoria County, New Brunswick.
E	Acadia Forest Experiment Station, Sunbury County, New Brunswick
F	St. Martins, St. John County, New Brunswick.
G	Oxford, Nova Scotia.

PLANTATIONS ESTABLISHED IN THE SPRING, 1961, FOR THE STUDY OF PREMATURE LOSS OF VIGOUR OF ABIES BALSAMEA Table 2.

			Nı	unber of	2 + 2 Tr	ees	
Origin of Seed	Newfou	ndland		New Bru	nswick		Nova Scotia
		_ل_ط	770	D	· El	Ω×,	Ċ
Acadia Forest Experiment Station, N. B.	576	576	1,2972/	1	8	* 1	i
Beaver Harbour, N. B.	I	i	I	ı	180	250	8
McGivney, N. B.	I	I	ł	I	4803/	009	60
Chisholm Brook, Green River, N. B.	I	I	ľ	100	420	2,200	50
Salmon River, St. John Co., N. B.	576	576	576	ſ	1,20	007	50
Avalon Peninsula, Nfld. (Seed lot MS.120-57	-	ı	ı	ı	8	1	ð
Burin Peninsula, Nfld.	ſ	8	I	ł	99	I	30
Sandy Brook, Nfld.	57641	/ 5764/	576	ì	300	100	I
Avalon Peninsula, Nfld. (Seed lot MS.232-57	1) 576	576	576	1	1	ł	100
Lake Melville, Labrador	ł	ı	J	I	8	I	J
Millerton District, Nfld.	I	I	I	1	30	8	1
	2,3044	2,3044	3,025	001	2,100	3,800	380
1/ Randomized blocks with four replications	3, 12 rov	ws of 1	2 trees	per plot	; spacing	g 61 x 61	

A A A 2/ Four plots of 144 trees each (576) plus 721 in double surround and single divis 3/ One plot of 90 trees plus double end rows and single division rows (390 trees) 4/ Plus division and surround trees.

0-7



Fig. 3. Locations of plantations in <u>Picea</u> abies provenance test

A and B	The Acadia Forest Experiment Station, Sunbury Co., N. B.
Cl and C2	Summit Depot, Green River Watershed, Restigouche Co., N. B.
Đ	Second Falls Reservoir, Madawaska County N. B.
E	University of New Brunswick, Fredericton, N. B.
F	Saint Martins, St. John County, N. B.
G	North Point, Queens County, Prince Edward Island.
H	East of Apple River, Cumberland County, N. S.
I	Near Advocate, Cumberland County, N. S.
J	Acadia University, Wolfville, N. S.
K	South of South Quinan, Yarmouth County, N. S.
L	Chignecto Game Sanctuary, Cumberland County, N. S.

PROVENANCE TEST	
ABIES	
PICEA	
II	
PLANTED	
TREES	
3	
+	
6F GF	
NUMBER	
Table 3.	

			Note Bran	de inter		1	• • •	н Сн С			Notes	Scotia		
			TO MONT	TATMOT										1
	А	р	C1.2/	c2 ^{2/}	Q	} E [⊞]	伍	сı	Н	н	Ŀ,	K	ц	
Picea abies														
Istebena, Poland	009	I	200	8	1000	200	001	1000	500	100	500	1000	t	
Nagold Schwarzwald, Germany	300	ł	200	8	200	200	007	500	I	100	500	100	570	
Westerhof, Harzen, Germany	300	ł	200	8	1000	200	001	500	500	100	500	500	1	
Thuringen, Germany	300	4	200	1	1000	200	00†	1000	500	001	500	500	9	
Schwabishen Alben, Germany	300	I	200	30	500	200	200	500	370	100	500	I	I	
Millstatt, Austria	300	ł	200	30	00,	200	00†	500	20	100	500	500	02	
Zwettl, Austria	300	I	200	8	001	200	ı	I	007	100	007	ł	P.	
Trieben, Austria	300	I	200	8	1000	200	1	I	001	100	500	1	ł	
Vopau, Austria	300	I	200	I	I	200	1	100	001	100	007	500	100	
Nauders, Austria	300	I	200	8	500	200	I	100	370	100	500	00†	07	
Kowary, Sudeten Mts., Poland	300	I	200	8	200	200	I	400	I	100	500	007	କ୍ଷ	
Halifax, N. S.	ł	500	200	30	I	200	1	007	300	100	007	007	1	1
Total P. abies:	3600	50	24,00	30	2300	24,00	2200	5600	1,240	1200	5700	4300	920	

٠

Table 3. (cont'd)

		New I	Brunswi	ck				P.E.I.			Nova S	scotia	
	₹¶	щ	c1 ^{2/}	C2 <u>5</u>	Q	E3/	ſ±.	ಅ	н	н	ſ	K	1 I
Picea glauca													
Acadia For. Exp. Sta., N. B.	12884	ł	200	ł	500	200	200	200	007	00T	I	200	1
Green River, N. B. (MS. 51-57)	1	I	200	t	500	200	I	500	500	100	t	007	1000
Green River, N. B. (MS.249-57)	ł	1	200	I	007	200	ı	007	400	100	T	400	1
Tobel P. glauca	1288	1	009	T	0071	009	200	0011	1300	300	- - -	1000	1000
Total trees:	4888	500	3000	300	8700	3000	2400	6700	5540	1-500	5700	5300	1900
, ,	51 ×61 (ótx61	6.61x	6.61x 6.61	~	1 9x1 9	61x61	81x81	61x6	उ [.] ह ट	//	181x81	81x81

 $\frac{1}{2}$ Design: 3 x 4 rectangular lattice with three replications and 100 trees per plot

2/ Cl - Established in spring 1961, with two replications of 100 trees per lot C2 - To be established in spring 1962

3/ Four replications of 50 trees per lot.

 $\frac{1}{2}/$ Division and surround trees

5/ Spacing: P. abies 8' x 8'; P. glauca 6' x 6'



Fig. 4. Locations of plantations of <u>Picea</u> <u>sitchensis</u>, <u>Picea</u> <u>glauca</u> and hybrids

Plantation	Location
A	The Acadia Forest Experiment Station, New Brunswick
B	Near Advocate, Nova Scotia.
C	Chignecto Game Sanctuary, Cumberland County, Nova Scotia.
D	Direct sowing at St. Martins, St. John County, New Brunswick, in 1957.

MATERIAL
GLAUCA
A I
AND
SITCHENSIS,
PICEA
5
DISTRIBUTION
Table 4.

its sent Petawawa 1060	750 7	0		0	0	0	0	0	0	0	0	•	009
Plai to 1	1777	200	200	R P	0	0	0	0	0	0	0	•	009
Seed Lots in Direct Sowing 1957	Ω		L L	Х Х м м	50 g	50 g				50 g			
þe	C	0		9 8 8	370	2300	0	0	0	2200	0	• •	5950
s Plant 1961	m	5	S S	01 01	100	100	0	0	0	OOL	100	ğ	800
Tree in	A	8	8 8		200	500	ц У	5	P P	80	500	<u>R</u>	2000
	Origin of Seed	Steward, Alaska	Holmer, Alaska	Vilsboel Vanned Plantage, Denmark	Lundsgaard, Denmark	Tornby Klit Plantage, Denmark	ST. <u>1</u> /179, Springfiel N. S.	Springfield, N. S.	Springfield, N. S.	Jutland, Denmark	Green River, N. B. (Seed lot MS.249-57)	Green River, N. B. (Seed lot MS.51-57)	
			(and x P. lutzii ?)	(and <u>P. sitchensis</u> x <u>glauca</u> ?)	8		2 − 3 22 − 3 22 − 3		P. glauca sitchensis ?)				Tree
	Species	sitchensis	, <mark>sitchensis</mark> sitchensis	sitchensis	2		2 •		. <u>glauca</u> (and x	glauca	£ ·	#	ST - Seed
	I	니	61 B	<u>'iei</u>	=	z ,	35 •	a +	ы	الم	=	#	ति

Few of the terminals had been browsed during the winter of 1960-1961. The terminal buds were again covered in October, 1961. At this time undesirable leaders were pruned, and dead and missing trees and the trees with no buds were replaced. The damage caused by browsing deer may complicate the interpretation of future measurements.

Table 5. SURVI	VAL OF PICEA RUBENS	Vean		- 1
Plantation ^{1/}	Location	Established	Percentage 1960	<u>survival²/</u> 1961
A	Central New Brunswick	1959	98.1 2 /	97.64
В	Northern New Brunswick	1960	N/A	98.5
C	Southern New Brunswick	1960	N/A	99•3

1/ Established according to 4 x 4 balanced lattice square designs with 16 seed sources, five replications, 100 trees per plot, division and surround rows and 4' x 4' spacing.

2/ Division and surround trees not included.

3/ No significant difference between survival of provenances (F value = 1.23)

4/ Survival since planting in 1959, dead and missing trees not replaced in 1960.

Plots of 100 trees per lot, from two lots of <u>Larix leptolepis</u> (Sieb. and Zucc.) Gord., five lots of <u>L. decidua</u> Mill. and one lot each of <u>L. decidua</u> <u>sudetica</u>, <u>L. decidua polonica</u> (Racib.) Ostenf. and Syrach, var., X <u>L. eurolepis</u> Henry and <u>L. laricina</u> (Du Roi) K. Koch, along with 50 <u>L. occidentalis</u> Nutt., 1150 trees in all, were planted at the Acadia Forest Experiment Station in the spring of 1961. About 700 trees, some from each lot were planted by the forestry students on the University of New Brunswick Woodlot. All lots, except the <u>L</u>. <u>laricina</u> came from the Petawawa Nursery. The seed source of each lot is known. An attempt to grow <u>Cedrus libani</u> Loud., <u>C. deodara</u> (Roxb.) Loud., <u>C. atlantica</u> Manetti and <u>C. atlantica glauca</u> Carr., resulted in complete mortality in all lots between sowing in the spring of 1958 and making estimates of nursery stock in 1960. <u>Larix leptolepis</u>, <u>L. decidua</u> and <u>L.</u> <u>laricina</u> for provenance studies will be outplanted in the spring of 1962. <u>Picea rubens</u> provenance material, and progenies of selected trees will be outplanted in 1963 and 1964 respectively.

As usual, plant material was exchanged with several research organizations.

REFERENCE

McNEEL, W., Jr., and J. KENNEDY. 1959. Prevention of browsing by deer in a pine plantation. J. Wildlife Mgmt., 23 (4): 450-451

CYTOGENETIC STUDIES IN CARAGANA

R.J. Moore

Plant Research Institute Research Branch, Canada Department of Agriculture, Ottawa

X-Irradiation of Seed

Dry seed of C. arborescens (open pollinated 1960 seed of several bushes mixed) was exposed to doses of 5,000r, 10,000r, 20,000r and 30,000r of X-irradiation in September 1960. Samples were germinated immediately and grown in greenhouse until the following May, when they were planted in the field.

Germination of seed of each treatment was 100%, of untreated seed, 62%. Irradiated seed began to germinate slightly (1-2 days) earlier than the control seed. There was no significant difference between the germination behaviour of seed of the different dosages.

Growth abnormalities were slight or absent in the seedlings of the 5,000r and 10,000r treatments but leaflets of the first 5-6 leaves of seedlings from the heavier treatments were crinkled, asymmetrical and mottled with yellow spots. Later leaves were normal and it appears that affected tissue did not survive competition with normal tissue. Root tips of the young seedlings were examined cytologically but no chromosome fragments were observed.

Measurements of the height of the plants in June 1961 indicated that seedlings from the 10,000r, 20,000r and 30,000r treatments were taller, by as much as 60%, than the controls but by the end of the growing season the controls equalled the tallest of the treated plants.

Some remaining seed of the 5,000r and 10,000r treatments and untreated seed was sown in October 1961 after storage for 1 year in envelopes under laboratory conditions. Germination was 100% (5,000r), 80% (10,000r), 66% (control). Leaf abnormalities were slight, as was the case in plants grown from seed of these treatments germinated immediately after germination. Storage did not seem to alter the irradiation affect.

Taxonomic Studies

As a result of taxonomic studies 2 papers were published:

Typification of the linnaean species of Caragana- Can. J. Botany 39, 1041-44 (1961). The type specimens of C. arborescens Lam., C. frutex (L.) Koch, C. pygmaea (L.) DC. were designated from those in the Linnaean Herbarium in London. The specimen bearing the epithet "Pygmaea" in Linnaeus' hand is actually C. frutex. It is proposed that another specimen, truly C. pygmaea, be designated as type to avoid renaming the species. It is apparent that Linnaeus understodd the species pygmaea and the error in labelling was merely a slip. On the origin of Caragana sinica \rightarrow J. Arnold Arb. 43, 203-214 (1962). Seed is distributed by many botanic gardens under the name C. sinica but it produces several other unrelated species. True C. sinica has been obtained only from the Grootendorst Nursery, Holland. This material is asexually propagated, is completely seed sterile and is triploid (2n-24). Pollen is partially aborted and normal embryo sacs are not formed. Herbarium specimens of the species from China have been examined, all have similar partially aborted pollen. On the basis of general morphology and pollen condition it is postulated that the species is a hybrid between C. rosea and C. microphylla. All available living material of both of the latter species is diploid and it is to be expected that their hybrid will be diploid and at least partially fertile. It is suggested that C. sinica in Asia is hybrid and diploid and that the triploid condition has arisen in cultivation through the formation of an unreduced gamete, the formation of which may have been stimulated by the unusual European environment.

REPORT TO COMMITTEE ON FOREST TREE BREEDING 1962.

bу

A.L. Orr-Ewing Research Division, B.C. Forest Service, Victoria, B.C.

1. General.

A graduate forester was added to the staff in October 1960. This increase in personnel has greatly benefited the research program. A new nursery area has been recently allocated. Deer-proof fences have been erected around the three outplanting areas at Lake Cowichan.

2. <u>Selection</u>.

The main emphasis has been on the selection of Douglas fir throughout Vancouver Island and the Lower Mainland. The objectives of this program are not only to select candidate trees for seed orchards but also to sample the population of coastal Douglas fir over as wide limits of its range as possible. Permanent clone banks of the trees selected are finally established for later breeding work. The number of Douglas fir selected from 1957 to 1961 is now 150; 64 of these trees being rated as plus and the remainder as almost plus trees. These trees are located over fifteen land districts in Vancouver Island and over two on the Lower Mainland. All of the trees selected to date have been propagated. The cruising involved has only been conducted during the summer months when temporary help was available so that progress has been slow and a great deal of the coastal region remains to be cruised. In addition, the work is becoming harder as the areas involved are becoming more and more inaccessible each year. Cooperation by industry is therefore, of the greatest assistance and is actively encouraged. Fortunately it has been increasing steadily since 1958, and culminated this year in the establishment of clone banks by four of the leading companies on the coast.

The establishment of the first Forest Service seed orchard has been purposely delayed until a sufficient number of suitable trees had been selected. It was considered advisable to have high clone numbers when the genetic quality of the material was completely unknown. The first seed orchard, therefore, is largely an experimental unit. The area for this orchard has now been cleared, fenced and planted to rootstock. The first propagations will be made in 1963.

3. Anatomical Characteristics of the Wood of Selected Trees.

Since 1958 the cores of 225 selected trees and their surrounding dominants have been sent to the Forest Products Laboratory in Vancouver. These cores were extracted with either a 5 mm, 12 mm or 3/4" borer. The

ALL BARRENE PLA PRIME DE LETTERE ALLE DE LETTERE

·. •

anger Stander og det Stander Stander Stander Stander Stander Stander Stander

್ರಾಂಗ್ಯಾಮಿ ಮಾಡಿ ಪ್ರವಾಸವರ್ಷವರ್ಷವರು ಮಾಡಿ ಸಂದಿನವರು ಮುಂದಾನವರು ಪ್ರವಾಸಿಗೆ ಬಿಡು ಆಟಿಯಲ್ಲಿ ಮಾಡಿದ್ದರು. ಪ್ರಾಟಿಕೆ ಸಂ 198 ರಿಸುದ್ರವರಿಗೆ ಸಂದ್ರಹಿಸಿ ಮುಂದು ಬಿರೆಗೆ ಸಂದ್ರಾ ಕಾಡಿಕೆ ಬರಿದಾರು ಪ್ರತಿದ ಮುಂದುಮಾಡಿಕ ಆಟಿಗಾದ್ದಾಗಿದ್ದು. ಬಿಟಿ 198 ರಿಸುದ್ರವರು ಬರಿಗೆ ಬಿರೆಗೆ ಬಿರೆಗೆ ಸಂದ್ರಾ ಬಿದ್ದಿ ಸಂದ್ರಾ ಬಿರೆಗೆ ಬರಿಗೆ ಮಾಡಿದ್ದಾಗೆ ಸಾಮಾನಕ ಸಹಿತಿ ಮಾಡಿದ್ದಾರೆ. ನಿರ್ಮ 198 ರಿಸುರ್ವಾರ್ ಸಿಂಗ್ರಾ ಮಾಡಿಕೆಗೆ ಬಿರೆಗೆ ಸ್ಥಾನ್ ಸೇವ್ ಸಂದ್ರಾ ಸಂದ್ರಾ ಸಿಂಗ್ರಾ ಮಾಡಿದ್ದಾರೆ. ಆಟಿಗಳು ಸಿಂಗ್ರಾ ಸಿಂಗ್ರಾ ಸಿಂಗ

and a second where a second stand we are not and the second second second second second second second second s and the second of the second a halas an an concentration and a sub-CAR 1990 TOP STA an arte at a constant and the second second and a second denses of the second at the second second second second and and construction of a construction of a construction of the construction of the construction of the second a single the set of the construction of the set of the Anodelia and contained and and a grant character of a second second burger and - Condition of the second of the second s I processes a set of the second sec niji Baranana analar kalamana nadaran batu yartuk butun butun butu (dalemenen barakbera lign two real with a size and a probability real capitor of the Probability of the part 《建建新路》),所以在白上在工作上书,所以为人,我的是一个个人的子子是这些新的人子。 计外国语 法非代理法 人名法法法德 第一次算法 We write we by pressing on a set of the state endogram the first of the set o systems and the set of the set of the set of the set of the

Figure 1. Experience of the number of each of the construction of the construction

s and the provide the second second

 specific gravity has been measured ring by ring from one to ten years on the cores of 64 of these trees and from eleven to thirty-five years at every five rings on 205 trees. Tracheid length has been measured in detail on 30 and spiral grain and compression wood on 205 trees. A complete report on this work is being prepared by the Forest Products Laboratory. It is hoped that this work will be continued as it is not only important that a background of information on the variation in internal characteristics of the Douglas fir be obtained, but that this investigation be extended to the progeny.

4. Progeny Tests.

A problem analysis involving the future of the whole genetic program has been recently written for detailed consideration by the Forest Service. It is essential that the most important aspect of the program, namely progeny testing, receives the necessary support.

The clone banks were all fertilised in the spring of 1961 and some interesting results were obtained. It was found that there was much variation in both cone and pollen production between different ramets of the same clone although all received the same amount of fertiliser. This was probably caused by the heterogeneity of the rootstock which, although from the same seed lot, would be highly variable as regards flowering propensities. Future research, therefore, will be directed towards the development of more uniform rootstock as irregular flowering within the ramets of a clone could have serious consequences in a seed orchard. Another clone was found to have a high proportion of hermaphroditic cones, a very undesirable character.

In general, the clones established from scions collected from the tops of sexually mature trees produced both cones and pollen in equal amounts. It was much more difficult, however, to induce pollen in young trees although cones could be produced in abundance in those as young as seven years. Further research into this problem of pollen induction is urgently needed. In spite of the shortage of pollen, some controlled crosses were possible in 1962.

5. Inbreeding of Douglas Fir.

The additional staff has enabled more time to be given to the breeding part of the program. Clones of some of the trees selected for the first seed orchard were inbred in 1962 to ascertain the degree of incompatibility. In 1961, successful backcrosses were made with four inbred progeny, in 1962 further backcrosses were made on a larger scale with progeny from five inbred lines. Sufficient pollen was also produced for some outcrosses to be made between different inbred lines, while further inbreeding to an S₂ generation was attempted with progeny in two lines.

6. <u>Selection in other Species.</u>

Western hemlock occurs in many of the stands which are cruised and trees are selected from time to time. As there is no great demand for the seed of this species for reforestation on the coast at the present time, the few trees selected have simply been propagated in a clone bank. M. Server S. S. Barrison, and the standard set of the standard set of the standard set of the partial standard set of the standard set of

and the second second

્યા ગામમાં આપણા આપણા ગામ પણ સમયમાં ગામ પ્રદુક્ષ્યના ગામ પ્રદાવ અને દોપ્ટ પ્રશિદ્ધ પ્રદાવ વૃદ્ધ સમયદ્ધા પ્રદાવ આપણે પાંચ ગામમાં ગામમાં ગામ આપણા સંદર્ભકાર્ટ પ્રસ્તાર દેવેલ દ્વાર્થક્રમના ઘટ્ટ દોપ્ય કે સમયદાર સંસ્થળદ્ધ કાર કર પિતાન પ્રધાર ગામમાં ગામમાં ગામમાં સાથાય કે સાથાય કે સાથે બાદ સંદર્ભના ઘટ્ટ સાંગ્રે કે સમયદાર વિસ્તાર દ્વાર કે શા આપણા સ્થાય કે બેઠા ગામમાં દુર્ભના સાથાય કે સાથાય કે સાથાય કે સાથે બંધ દોપ્ય ગામ સાથાય કે સાથ પ્રદાવ સાથાય કે સાથ

Second and the set of a start all for the set of the spirit print of the second set of the second s

Es marat sit and solves even as a sex if and (these posture) (the solves) summer is if the sex if and (these posture) (the solves) and poster in the second state if the sex if a second solves) is a second solver. A second solves as a second solves is a second solves of the second solves in the second solves. It is a second solves of the second solves in the second solves. It is a second solves of the second solves in the second solves in the second solves of the second solves in the second solves in the second solves. It is a second solves of the second solves of the second solves in the second solves in the second solves of the second solves in the second solves of the second solves in the se

and the first a the state of

A base of the solution of the solutio

(a)
 (b)
 (c)
 <li(c)
 <li(c)
 <li(c)
 (c)

contracted office of the contraction of

are entrypastang and the theory of the subsection of the control of a constant of the conare anticed to the relation to the end of the constant of the subsection the constant of the first constant to the relation of the constant of the best taken the constant of the constant is constant to the constant of the constant of the best taken of the constant of the second to the constant of the constant of the second of the constant of the constant of the second of the constant of the second of the constant of the constant of the second of the constant of t

7. Plantings of Exotic Conifers for Future Hybridization.

.

The small breeding arboretum started in 1960 has now been restricted to the genera <u>Pseudotsuga</u> and <u>Tsuga</u> only. All the known species of these two genera are now represented in the arboretum at Lake Cowichan. Seed of many other species is sent to the recently established arboretum at the University Forest of the University of British Columbia.

. .

The space control of the trace of the matrice of the matrix and the control the trace of the space of the second to the trace of t

"SUMMARY REPORT" 1961-62 par L. PARROT Faculté d'Arpentage et de Génie Forestier Université Laval Québec, P. Qué.

I. Etude de Provenances.

10 Une plantation de 25 provenances de pin rouge a été mise sur pied suivant dispositif statistique avec répétitions, en collaboration avec la station de Recherches Forestières de Petawawa, Ont.

20 Les semences de 95 provenances de pin gris, couvrant l'aire canadienne, ont été ensemencées en mai 1962. Etude effectuée avec l'organisme de recherches ci-haut mentionné.

30 Quelque dix races de pin sylvestre, provenant de tiges sélectionnées et de vergers à graines, ont été ensemencées simultanément avec 20.

40 Plantation de quelque 115 provenances de pin sylvestre, qui se continuera en 1963, en collaboration avec le professeur J. W. Wright, de l'Université du Michigan, E. U.

50 Une plantation de clones de <u>Populus</u>, sp. a été amorcée et se continuera incessamment.

II. Sélection individuelle.

La sélection d'arbres plus de Pin rouge et d'épinette noire est commencée depuis 1961 et portera sur plusieurs années.

III. Enseignement.

Un cours nouveau et couvrant les différents aspects de la génétique forestière et de l'amélioration des arbres forestiers est donné à la Faculté; ce dernier étant surtout orienté vers un but pratique. IV. Publications.

1) "L'importance de la sélection en reboisement", Forêt Conservation, mars 1962.

2)"La génétique forestière, une clé pratique au reboisement" Le Laval forestier et géomètre, mai 1962.

V. Travaux en perspective.

1) Etude sur un aspect génétique de l'Acer saccharum.

2) Approche sur la floraison de certains conifères d'âge juvénile.

LP/SDV

REPORT TO COMMITTIL ON FOREST TREE BRIEDING, AUGUST, 1962.

White Pine Meevil: Tree Resistance Studies C. R. Sullivan Forest Insect Laboratory, Box 490, Sault Ste. Marie, Ontario.

I. Introduction

The fact that adults of the white pine weevil, <u>Pissodes</u> <u>strobi</u> Peck, have a definite preference for white pine leaders of particular quality led Dr. C. Heimburger to study the resistant qualities of pine stock, selected specifically for good growth, form, and seemingly resistance to weetil attack, under natural conditions. The studies were begun during the spring of 1957, at the Kirkwood Management Unit, Ontario Department of Lands and Forests, Thessalon, Ontario. In co-operation with Dr. Heimburger, the Ferest Insect Laboratory, Sault Ste. Marie, has assumed responsibility for inspecting test grafts set out in field experiments to obtain records on the success, growth characteristics, and incidence of weevilling in the selected stock.

II. Experiment No. 1

This experiment involved grafting into pruned Scots pine, 400 scions made up of 20 grafts from each of five clones of resistant <u>P. strobus</u> and five clones of susceptible <u>P. strobus</u>, plus 20 grafts from each of five clones of resistant <u>P. peuce</u> and susceptible <u>P. peuce</u>. Annual surveys during the period 1958 to 1961 revealed considerable variability in their survival and vigor. The <u>P. peuce</u> grafts should a higher percentage of survival than did the <u>P. strobus</u> grafts, but among the survivors of both species graft vigor has been about equal, with about 60 per cent of the grafts showing good to moderate vigor, and the remaining 40 per cent showing poor vigor.

Analysis of weevilling has been limited to stoms of moderate to good vigor, because it is typical of white pine weevil adults to confine their attack to leaders so characterized. Of the resistant <u>P. strobus</u> stock, none of the leaders were killed as a result of weevil attack in 1959, 1960, or 1961, whereas among the susceptible stock, five, three, and 15 per cent of the leaders were killed, respectively. Of the resistant <u>P. peuce</u> stock, one, five, and five per cent of the leaders were killed by the weevil in 1959, 1960, and 1961, respectively, whereas among the susceptible stock, four 16, and 14 per cent were killed respectively. These figures indicate that both <u>P. strobus</u> and <u>P. peuce</u> stock, assumed to be susceptible to weevil attack at the beginning of the experiment, are indeed, more susceptible than the stock classified as capable of resisting weevil attack.

Tests of resistance of the above stock to blister rust were carried out by Dr. Heimburger concurrently with these studies. His results indicate that the <u>P. strobus</u> stock used in this experiment is rust susceptible, but that the <u>P. peuce</u> clones, especially those that seem to be somewhat more resistant to weevil attack, merit further consideration in the future. Examinations of this stock will, therefore, be continued. III. <u>Experiment No. 2</u>

In 1961, the scope of the project was increased to include an additional 1,050 white pine trees top-grafted with 50 scions of each of 21 clones collected in Canada and United States. Materials grafted include <u>P. strobus</u>, <u>P. peuce</u>, <u>P</u>.

U-2

<u>griffithil</u>, and <u>P. monticola</u>, in addition to hybrids of each of the last three species obtained from crossings with <u>P.</u> <u>strobus</u>, and hybrids obtained from crossing <u>P. strobus</u> with <u>Pi</u> <u>parviflora</u>. The grafts were selected for resistance to blister rust and resistance or susceptibility to weevil attack. During 1961, chemicals were used to protect the grafts from weevil attack and permit maximum graft success. Eighty-eight per cent of the grafts from the 21 clones (range between clones, 50 to 100 per cent) survived the summer and about 90 per cent of these exhibited good vigor. It is not known at this time whether grafts of the various clones will be fully hardy in the Kirkwood area. Observations on this aspect of the problem are continuing.

シート・ション (1993年) (1993年) (1993年) (1993年) (1993年)

and the second secon

A ment of the second s second sec

and the second of the production of the second s

Ale Alexandra and <u>alexandra alexandra alexandra alexandra alexandra alexandra alexandra alexandra alexandra ale</u> **by:**

Oscar Sziklai

- State Added and a state of the University of British Columbia & Wetter and the University of Forestry all and the Added and the Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and the Added and Faculty of Forestry all and the Added and Faculty of Forestry and the Faculty of Forestry and the Faculty of Faculty of

(a) Experience and the section of the second secon second sec

Report on Tree Improvement Work at

The University of British Columbia, Faculty of Forestry

Forest Tree Improvement work at the University of British Columbia is chiefly concerned with Douglas fir, the most important timber species on the Pacific Coast. Western hemlock, western red cedar, lodgepole pine, several native Abies species, black cottonwood, and some hybrid poplars are also under study.

The following projects were active during 1960 and 1961.

1. <u>Studies of Douglas fir provenances</u>, with special reference to British Columbia.

Investigators: Dr. P. G. Haddock and J. Walters.

The objective is to supplement studies under way in coastal Douglas fir provenances, by the Oregon Forest Lands Research Center. Thirteen provenances were included, mostly from the Interior regions of the Province. The seedlings were observed in the first two years in the nursery bed. Phenological observation and growth characteristics were recorded and 2+0 seedlings were transplanted to the University Research Forest in the spring of 1961. Further observations on growth characteristics are continuing.

 The effect of inheritance on germination behavior of seed of Douglas fir. Initiated by Dr. G. S. Allen, continued by O. Sziklai.

The objective is the determination of the physiological basis for dormancy and the relation of this to inheritance.

Intraspecific reciprocal crosses were carried out on three trees in 1958, 1959 and 1961. Dry and wet pollination methods were tried. The wet pollination method gave double the number of viable seed produced by the dry pollination method. Stratified and unstratified seeds were tested by standard methods for germination. Seedlings were transplanted in 1959 to the University Campus Nursery. Progeny tests will be established at the University Research Forest in 1963 for later work with the F_2 generation and back crosses.

. · · .

3. Nutrition of seedlings of Douglas fir.

Investigators: Dr. G. S. Allen, Dr. D. J. Wort and Dr. J. H. G. Smith. The objective is the development of better planting stock and the determination of nutritional requirements of Douglas fir.

Two runs have been made in the nursery; one with inorganic amendments only and the other with organic amendments from specific forest site-types. The experiment was repeated in 1960 and supplemented with 38 seedlots collected from individual trees. Growth characteristics were obtained from the parent trees and seedling characteristics were correlated with the parent characteristics.

4. <u>The effect of release from competition on certain wood characteristics of</u> Douglas fir and western hemlock.

Investigator: Dr. R. W. Wellwood.

.

Increment cores (3/4-in.dia.) were obtained from 72 trees released by a 1949 thinning in a 70-year-old Douglas fir-hemlock stand. Sampling included four site indices and three crown classes. Core samples were tested for changes in rate of growth, specific gravity, fiber length and fibril angle in the growth periods 1944-49 and 1950-55.

5. The selection and testing of Douglas fir and western hemlock plus trees.

Investigators: Dr. R. W. Wellwood, Dr. P. G. Haddock and J. Walters. The objective is to locate and propagate superior phenotypes of Douglas fir and western hemlock. 2250 acres were reconnoitred and 812 acres were cruised 100 per cent on the University Research Forest in 1959. Field data were collected on 16 Douglas fir and on 20 western hemlock trees, which included: age, height, d.b.h., growth rate, stem form, crown form, length of live crown, branching habit, branch size, bark type and thickness, cone production, resistance to insect and disease. Sites were described, and photographs were taken.

Wood samples were collected for laboratory analysis; fiber length, specific gravity, fibril angle, sapwood-heartwood ratio, springwood-summerwood ratio, cellulose content were defined.

On the basis of the data collected the candidate trees were classified as volume-plus, form-plus or quality-plus trees.

V-3
5. Propagation of the selected trees by means of rooting and grafting is in progress.

V-4

6. <u>Selection and testing of variation among poplar species</u>. Investigators: Dr. J. H. G. Smith, J. Walters and G. Blom. Native cottonwood from several sources are being compared with the hybrid poplars <u>Populus grandis</u>, <u>Populus robusta</u> and <u>Populus regenerata</u> on a

commercial scale on about 600 acres of plantations. Populations containing 69 of the Oxford hybrid poplars, 10 Italian hybrid

poplars, and a limited number of native black cottonwoods were established at the University Research Forest and on the Harper property near Agassiz.

Selection and testing of variation among individual black cottonwoods.
Investigators: Dr. J. H. G. Smith and Dr. R. W. Kennedy.

Fifteen trees were chosen on the basis of superior growth rate and form and were grown in a nursery. Measurements of fiber length and specific gravity of some of these were made to assess the relative importance of inherited and environmental characteristics.

 <u>Natural variations in four species of Abies from British Columbia</u>. Investigator: K. J. Roller.

Twigs and cones were collected from 29 populations of the four native species of the Province, through the range of six degrees of latitude and 7,000 feet in elevation.

Analysis is being made of the taxonomic variation in needle sections, in cone scales and bracts in order to determine the standard and variable morphological and anatomical characteristics of these species. Additional studies have been made of germination and of division of roottip cells.

9. <u>Variation in lodgepole pine, and its role in the genetic improvement of</u> <u>Coastal forms</u>.

Investigator: L. Roche.

las sub-

The growth habit of the species under natural conditions was observed in the field. Growth habit and growth rates were studied.

9. Leaf measurements and germination tests were made on specimens of foliage and seed collected throughout the distribution of the species. Variation in bark type and color was studied in the field and was recorded by means of photographs.

Recommendations were made for future selection and breeding work.

10. Study of individual tree variations of Douglas fir.

Investigator: Dr. B. G. Griffith.

The objective is to explore the intraspecific variation of Douglas fir and to determine whether these characteristics are inherited or controlled by environments.

A total of 155 open-grown young Douglas fir trees were selected at the University Research Forest in 1956. Age, height, d.b.h., crown length and width and growth rate were described. Phenological observations on radial and vertical growth, flowering characteristics, cone production were observed from 1956 every year.

Controlled crosses were attempted by O. Sziklai in 1962 and will be carried out in the later years to determine how the different characteristics are inherited. These trees also are being studied by A. Kozak to determine factors governing the incidence of attack by cone and seed insects.

11. <u>Analysis of variation of Douglas fir, western hemlock, and western red</u> <u>cedar</u>.

Investigators: Dr. J. H. G. Smith, Dr. J. W. Ker, and J. Csizmazia. In order to make a preliminary assessment of the possibilities of improvement of tree and wood quality by selection and breeding, data on variation in wood and tree characteristics were colk cted and analyzed. Natural pruning was studied intensively and factors governing tree and branch size analyzed thoroughly. The importance of measurement of crown width to determine the influence of competition on invidivual trees as a guide to desired spacing was stressed. It is essential that in selection of individual trees for study due consideration be given to their competitive status which can be expressed conveniently by the ratio of crown width to d.b.h.

12. A program for improvement of basket willow on the Carpathian Plain of Hungary. and a substance to make a second as the (e.g., Investigator: . O. Sziklaipe et egy arder for the difference An inventory of the most important indigenous and exotic Salix clones for basket making was carried out in Hungary. Eleven clones were selected Agene from those presently growing in the various parts of the country. The vegetative and generative characteristics of the shoots were described. Interspecific crosses were carried out using five different combinations. The germination of Salix seed was studied by taking a series of photographs. kan ya € ing i Five experimental plantations were set out in the more important soil regions, and the eleven clones were planted in randomized blocks. Height measurements were recorded bi-weekly during the growing season, and more than 50,000 were made each year from 1955 to 1959. The influence of the clone, year, and locality on the average height en an Sterre attained was evaluated by means of an analysis of variance, and components 4.5.2 (20) of variance were used to estimate the heritability. Simple correlation coefficients were calculated to describe degree of association of average 4度が見たり、 「「」」」。 height, weight and volume of different clones in the five localities during the years 1955-59 inclusive. S. and Sugar 18 1 B B B - 1 B -13. General analyses of chemical, anatomical, and physical properties of wood. Investigators: Dr. R. W. Wellwood, Dr. R. W. Kennedy in i par A large number of individual projects have been undertaken or are under way to determine factors variation in wood quality and factors that govern it. in the second state of the second states in the second states of the second states and the second states are se 14. Theses for the B.S.F. degree. 1 1 1 1 2 2 Investigators: Senior students and staff, Each year several senior theses provide new information on problems in the field of forest genetics but these are not listed here. 15. Projects on the University Research Forest.

Investigators: J. Walters and J. Spos.

Projects in silvics and silviculture of west coast species studied at the University Research Forest are reported upon annually but have not been listed here.

V-6

Publications

- 1. Allen, G.S. 1960. A method of distinguishing Coastal from Interior Douglas fir seed. British Columbia Lumberman. 44: 26-30.
- 2. Allen, G.S. 1961. Testing Douglas fir seed for provenance. Proceedings of International Seed Testing Association. Vol.26.No.3.pp.308-403.
- 3. Allen, G.S. and O. Sziklai. 1962. Pollination of Douglas-fir with water suspensions of pollen. Forest Science. Vol.8, No.1.
- Kennedy, R.W. and J.M. Jaworsky. 1960. Variation in cellulose content of Douglas fir. Tappi 43: 25-27.
- 5. Roche, L. 1962. Variation in <u>Pinus contorta</u>, and its role in the genetic improvement of Coastal forms. Master of Forestry thesis. Faculty of Forestry, U.B.C.
- 6. Smith, J.H.G., J.W. Ker, and J. Csizmazia. 1961. Economics of reforestation of Douglas fir, western hemlock and western red cedar in the Vancouver Forest District. Faculty of Forestry Bulletin No. 3.
- 7. Sziklai, O. 1961. A program for improvement of basket willow on the Carpathian Plain of Hungary, Master of Forestry thesis. Faculty of Forestry, U.B.C.
- Walters, J., J. Soos and P.G. Haddock. 1961. The selection of plus trees on the University of British Columbia Research Forest, Haney, B. C. Research Papers, Faculty of Forestry U.B.C. No.33.
- 9. Wellwood, R.W. and J.H.G. Smith, 1962. Variation in some important qualities of wood from young Douglas fir and western hemlock trees. Research Papers, Faculty of Forestry, U.B.C. No.50.